

SUBSTITUTED INDOLINES WHICH INHIBIT RECEPTOR TYROSINE KINASES

Related Applications


- 5 Benefit of U.S. Provisional Application Serial No. 60/160,547, filed on October 20, 1999, is hereby claimed.

Field of the Invention

- 10 The present invention relates to novel indolinones that inhibit receptor tyrosine kinases, their use as pharmaceuticals, particularly in the treatment of proliferative diseases, and pharmaceutical compositions comprising these compounds.

Description of the Invention

- 15 The present invention provides new indolinones of general formula



- 20 substituted in the 6 position, the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof, particularly the physiologically acceptable salts thereof which have valuable properties.

- The above compounds of general formula I wherein R₁ denotes a hydrogen atom or a prodrug group have valuable pharmacological properties, in particular an inhibiting effect on various kinases, especially receptor tyrosine kinases such as VEGFR2, PDGFR α , PDGFR β , FGFR1, FGFR3, EGFR, HER2, IGF1R and HGFR, as well as complexes of
- 25

CDK's (Cyclin Dependent Kinases) such as CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK8 and CDK9 with their specific cyclins (A, B1, B2, C, D1, D2, D3, E, F, G1, G2, H, I and K) and to viral cyclin (cf. L. Mengtao in J. Virology 71(3), 1984-1991 (1997)), and on the proliferation of cultivated human cells, in particular endothelial cells,
5 e.g. in angiogenesis, but also on the proliferation of other cells, in particular tumour cells.

The other compounds of the above general formula I wherein R₁ does not denote a hydrogen atom or a prodrug group are valuable intermediate products for preparing the
10 abovementioned compounds.

The present invention thus relates to the above compounds of general formula I, whereby those compounds wherein R₁ denotes a hydrogen atom or a prodrug group have valuable pharmacological properties, pharmaceutical compositions containing the
15 pharmacologically active compounds, the use thereof and processes for preparing them.

In the above general formula I

20 X denotes an oxygen or sulphur atom,

R₁ denotes a hydrogen atom or a prodrug group such as a C₁₋₄-alkoxycarbonyl or C₂₋₄-alkanoyl group,

25 R₂ denotes a carboxy group, a straight-chain or branched C₁₋₆-alkoxy-carbonyl group, a C₄₋₇-cycloalkoxy-carbonyl or an aryloxycarbonyl group,

a straight-chain or branched C₁₋₆-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, heteroaryl, carboxy, C₁₋₃-alkoxy-carbonyl, aminocarbonyl,
30 C₁₋₃-alkylamino-carbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group,

a straight-chain or branched C₂₋₆-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a chlorine atom or a hydroxy, C₁₋₃-alkoxy, amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group,

- 5 an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C₁₋₃-alkoxy group or, if R₄ does not denote an aminosulphonyl-phenyl or N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl-phenyl group, it may also denote a di-(C₁₋₂-alkyl)-aminocarbonyl group,

- 10 R₃ denotes a hydrogen atom, a C₁₋₆-alkyl, C₃₋₇-cycloalkyl, trifluoromethyl or heteroaryl group,

- a phenyl or naphthyl group, a phenyl or naphthyl group mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a trifluoromethyl, C₁₋₃-alkyl or C₁₋₃-alkoxy group, whilst in the event of disubstitution the substituents may be identical or different and wherein the abovementioned unsubstituted as well as the mono- and disubstituted phenyl and naphthyl groups may additionally be substituted

- by a hydroxy, hydroxy-C₁₋₃-alkyl or C₁₋₃-alkoxy-C₁₋₃-alkyl group,

- 20 by a cyano, carboxy, carboxy-C₁₋₃-alkyl, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylamino-carbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group,

- by a nitro group,

- 25 by an amino, C₁₋₃-alkylamino, di-(C₁₋₃-alkyl)-amino or amino-C₁₋₃-alkyl group,

- by a C₁₋₃-alkylcarbonylamino, N-(C₁₋₃-alkyl)-C₁₋₃-alkyl-carbonylamino, C₁₋₃-alkylcarbonylamino-C₁₋₃-alkyl, N-(C₁₋₃-alkyl)-
30 C₁₋₃-alkylcarbonylamino-C₁₋₃-alkyl, C₁₋₃-alkyl-sulphonylamino,

C₁₋₃-alkylsulphonylamino-C₁₋₃-alkyl, N-(C₁₋₃-alkyl)-
C₁₋₃-alkylsulphonylamino-C₁₋₃-alkyl or aryl-C₁₋₃-alkylsulphonylamino group,

by a cycloalkylamino, cycloalkyleneimino, cycloalkyleneiminocarbonyl,
cycloalkyleneimino-C₁₋₃-alkyl, cycloalkyleneiminocarbonyl-C₁₋₃-alkyl or
cycloalkyleneiminosulphonyl-C₁₋₃-alkyl group having 4 to 7 ring members in each
case, whilst in each case the methylene group in position 4 of a 6- or 7-membered
cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a
sulphinyl, sulphonyl, -NH or -N(C₁₋₃-alkyl) group,

or by a heteroaryl or heteroaryl-C₁₋₃-alkyl group,

R₄ denotes a C₃₋₇-cycloalkyl group,


whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group
may be substituted by an amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group or
replaced by an -NH or -N(C₁₋₃-alkyl) group,

or a phenyl group substituted by the group R₆, which may additionally be mono- or
disubstituted by fluorine, chlorine, bromine or iodine atoms, by C₁₋₅-alkyl, trifluoromethyl,
hydroxy, C₁₋₃-alkoxy, carboxy, C₁₋₃-alkoxycarbonyl, amino, acetylamino,
C₁₋₃-alkyl-sulphonylamino, aminocarbonyl, C₁₋₃-alkyl-aminocarbonyl, di-(C₁₋₃-al-
kyl)-aminocarbonyl, aminosulphonyl, C₁₋₃-alkyl-aminosulphonyl,
di-(C₁₋₃-alkyl)-aminosulphonyl, nitro or cyano groups, wherein the substituents may be
identical or different and wherein

R₆ denotes a hydrogen, fluorine, chlorine, bromine or iodine atom,

a cyano, nitro, amino, C₁₋₅-alkyl, C₃₋₇-cycloalkyl, trifluoromethyl, phenyl, tetrazolyl or
heteroaryl group,

the group of formula



wherein the hydrogen atoms bound to a nitrogen atom may in each case be replaced independently of one another by a C₁₋₃-alkyl group,

5

a C₁₋₃-alkoxy group, a C₁₋₃-alkoxy-C₁₋₃-alkoxy, phenyl-C₁₋₃-alkoxy, amino-C₂₋₃-alkoxy, C₁₋₃-alkylamino-C₂₋₃-alkoxy, di-(C₁₋₃-alkyl)-amino-C₂₋₃-alkoxy, phenyl-C₁₋₃-alkylamino-C₂₋₃-alkoxy, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino-C₂₋₃-alkoxy, C₅₋₇-cycloalkyleneimino-C₂₋₃-alkoxy or C₁₋₃-alkylmercapto group,

10

a carboxy, C₁₋₄-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylamino-carbonyl, N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl, phenyl-C₁₋₃-alkylamino-carbonyl, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino-carbonyl, piperazinocarbonyl or N-(C₁₋₃-alkyl)-piperazinocarbonyl group,

15

a C₁₋₃-alkylaminocarbonyl or N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl group wherein an alkyl moiety is substituted by a carboxy or C₁₋₃-alkoxycarbonyl group or in the 2 or 3 position by a di-(C₁₋₃-alkyl)-amino, piperazino, N-(C₁₋₃-alkyl)-piperazino or a 4- to 7-membered cycloalkyleneimino group,

20

a C₃₋₇-cycloalkyl-carbonyl group,

wherein the methylene group in the 4 position of the 6- or 7-membered cycloalkyl moiety may be substituted by an amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group or replaced by an -NH or -N(C₁₋₃-alkyl) group,

25

a 4- to 7-membered cycloalkyleneimino group wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

the cycloalkylene moiety may be fused to a phenyl ring or

5

one or two hydrogen atoms may each be replaced by a C₁₋₃-alkyl group and/or

10

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, phenyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino group or

15

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl), -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group,

a C₁₋₄-alkyl group substituted by the group R₇, wherein

R₇ denotes a C₃₋₇-cycloalkyl group,

20

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group or replaced by an -NH or -N(C₁₋₃-alkyl) group or

25

in a 5- to 7-membered cycloalkyl group a -(CH₂)₂ group may be replaced by a -CO-NH group, a -(CH₂)₃ group may be replaced by a -NH-CO-NH or -CO-NH-CO group or a -(CH₂)₄ group may be replaced by a -NH-CO-NH-CO group, whilst in each case a hydrogen atom bound to a nitrogen atom may be replaced by a C₁₋₃-alkyl group,

30

an aryl or heteroaryl group,

a hydroxy or C₁₋₃-alkoxy group,

an amino, C₁₋₇-alkylamino, di-(C₁₋₇-alkyl)-amino, phenylamino,
N-phenyl-C₁₋₃-alkyl-amino, phenyl-C₁₋₃-alkylamino, N-(C₁₋₃-alkyl)-phenyl-
C₁₋₃-alkylamino or di-(phenyl-C₁₋₃-alkyl)-amino group,

an ω-hydroxy-C₂₋₃-alkyl-amino, N-(C₁₋₃-alkyl)-ω-hydroxy-C₂₋₃-alkyl-amino,
di-(ω-hydroxy-C₂₋₃-alkyl)-amino, di-(ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl)-amino or N-(dioxo-
lan-2-yl)-C₁₋₃-alkyl-amino group,

a C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-amino or
C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-N-(C₁₋₃-alkyl)-amino group,

a C₁₋₃-alkylsulphonylamino, N-(C₁₋₃-alkyl)-C₁₋₃-alkylsulphonylamino,
C₁₋₃-alkylsulphonylamino-C₂₋₃-alkyl-amino or
C₁₋₃-alkylsulphonylamino-C₂₋₃-alkyl-N-(C₁₋₃-alkyl)-amino group,

a hydroxycarbonyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-
hydroxycarbonyl-C₁₋₃-alkyl-amino group,

a guanidino group wherein one or two hydrogen atoms may each be replaced by a C₁₋₃-alkyl group,

a group of formula




wherein

R₈ denotes a hydrogen atom or a C₁₋₃-alkyl group,

n denotes one of the numbers 0, 1, 2 or 3 and

R₉ denotes an amino, C₁₋₄-alkylamino, di-(C₁₋₄-alkyl)-amino, phenylamino, N-(C₁₋₄-alkyl)-phenylamino, benzylamino, N-(C₁₋₄-alkyl)-benzylamino or C₁₋₄-alkoxy group, a 4- to 7-membered cycloalkyleneimino group, whilst in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl), -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group, or, if n denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula



wherein

R₁₀ denotes a hydrogen atom, a C₁₋₃-alkyl group, a C₁₋₃-alkylcarbonyl, arylcarbonyl, phenyl-C₁₋₃-alkyl-carbonyl, C₁₋₃-alkylsulphonyl, arylsulphonyl or phenyl-C₁₋₃-alkylsulphonyl group,

m denotes one of the numbers 1, 2, 3 or 4,

o denotes the number 1 or, if m denotes one of the numbers 2, 3 or 4, o may also denote the number 0 and

R₁₁ denotes an amino, C₁₋₄-alkylamino, di-(C₁₋₄-alkyl)-amino, phenylamino, N-(C₁₋₄-alkyl)-phenylamino, benzylamino, N-(C₁₋₄-alkyl)-benzylamino, C₁₋₄-alkoxy or C₁₋₃-alkoxy-C₁₋₃-alkoxy group, a di-(C₁₋₄-alkyl)-amino-C₁₋₃-alkylamino group optionally substituted in the 1 position by a C₁₋₃-alkyl group or a 4- to 7-membered cycloalkyleneimino group, wherein the cycloalkylene

moiety may be fused to a phenyl ring or in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl), -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group,

5

a C₄₋₇-cycloalkylamino, C₄₋₇-cycloalkyl-C₁₋₃-alkylamino or C₄₋₇-cycloalkenylamino group wherein position 1 of the ring is not involved in the double bond and wherein the abovementioned groups may each additionally be substituted at the amino-nitrogen atom by a C₅₋₇-cycloalkyl, C₂₋₄-alkenyl or C₁₋₄-alkyl group,

10

a 4- to 7-membered cycloalkyleneimino group, wherein

the cycloalkylene moiety may be fused to a phenyl group or to an oxazolo, imidazolo, thiazolo, pyridino, pyrazino or pyrimidino group optionally substituted by a fluorine, chlorine, bromine or iodine atom, by a nitro, C₁₋₃-alkyl, C₁₋₃-alkoxy or amino group, and/or

15

one or two hydrogen atoms may each be replaced by a C₁₋₃-alkyl, C₅₋₇-cycloalkyl or phenyl group and/or

20

the methylene group in the 3 position of a 5-membered cycloalkyleneimino group may be substituted by a hydroxy, hydroxy-C₁₋₃-alkyl, C₁₋₃-alkoxy or C₁₋₃-alkoxy-C₁₋₃-alkyl group,

25

the methylene group in the 3 or 4 position of a 6- or 7-membered cycloalkyleneimino group may in each case be substituted by a hydroxy, hydroxy-C₁₋₃-alkyl, C₁₋₃-alkoxy, C₁₋₃-alkoxy-C₁₋₃-alkyl, carboxy, C₁₋₄-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, phenyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkyl-amino group or

30

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl-), -N(phenyl), -N(phenyl-C₁₋₃-alkyl-), -N(C₁₋₃-alkyl-carbonyl-), -N(C₁₋₄-hydroxy-carbonyl-), -N(C₁₋₄-alkoxy-carbonyl-), -N(benzoyl-) or -N(phenyl-C₁₋₃-alkyl-carbonyl-) group,

5

wherein a methylene group linked to an imino-nitrogen atom of the cycloalkyleneimino group may be replaced by a carbonyl or sulphonyl group or in a 5- to 7-membered monocyclic cycloalkyleneimino group or a cycloalkyleneimino group fused to a phenyl group the two methylene groups linked to the imino-nitrogen atom may each be replaced by a carbonyl group,

10


or R₆ denotes a C₁₋₄-alkyl group which is substituted by a carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group or by a 4- to 7-membered cycloalkyleneiminocarbonyl group,

15

an N-(C₁₋₃-alkyl)-C₂₋₄-alkanoylamino group which is additionally substituted in the alkyl moiety by a carboxy or C₁₋₃-alkoxycarbonyl group,

20

a group of formula



25 wherein

R₁₂ denotes a hydrogen atom, a C₁₋₆-alkyl or C₃₋₇-cycloalkyl group or a C₁₋₃-alkyl group terminally substituted by a phenyl, heteroaryl, trifluoromethyl, hydroxy, C₁₋₃-alkoxy, aminocarbonyl, C₁₋₄-alkylamino-carbonyl, di-(C₁₋₄-alkyl)-amino-carbonyl, C₁₋₃-alkyl-carbonyl, C₁₋₃-alkyl-sulphonylamino,

30

N-(C₁₋₃-alkyl)-C₁₋₃-alkyl-sulphonylamino, C₁₋₃-alkyl-aminosulphonyl or di-(C₁₋₃-alkyl)-aminosulphonyl group and


p denotes one of the numbers 0, 1, 2 or 3 and

5

R₁₃ assumes the meanings of the abovementioned group R₇, or, if p denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

10



wherein

15

R₁₄ denotes a hydrogen atom, a C₁₋₄-alkyl group, a C₁₋₃-alkylcarbonyl, arylcarbonyl, phenyl-C₁₋₃-alkylcarbonyl, heteroarylcarbonyl, heteroaryl-C₁₋₃-alkylcarbonyl, C₁₋₄-alkylsulphonyl, arylsulphonyl, phenyl-C₁₋₃-alkylsulphonyl, heteroarylsulphonyl or heteroaryl-C₁₋₃-alkyl-sulphonyl group,

20

q denotes one of the numbers 1, 2, 3 or 4,


r denotes the number 1 or, if q is one of the numbers 2, 3 or 4, it may also denote the number 0 and

25

R₁₅ assumes the meanings of the abovementioned group R₇,

a group of formula

30



wherein

R₁₆ denotes a hydrogen atom or a C₁₋₄-alkyl group optionally terminally substituted by a cyano, trifluoromethyl-carbonylamino or N-(C₁₋₃-alkyl)-trifluoromethyl-carbonyl-amino group and

5

R₁₇ denotes a C₁₋₃-alkyl group,

an amino group substituted by a di-(C₁₋₃-alkyl)-amino-C₁₋₃-alkyl-carbonyl or di-(C₁₋₃-alkyl)-amino-C₁₋₃-alkyl-sulphonyl group and a di-(C₁₋₃-alkyl)-aminocarbonyl-C₁₋₃-alkyl group,

10

or an N-(C₁₋₃-alkyl)-C₁₋₅-alkylsulphonylamino or N-(C₁₋₃-alkyl)-phenylsulphonylamino group wherein the alkyl moiety is additionally substituted by a cyano or carboxy group,

15

wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R₆ may be mono- or disubstituted by fluorine, chlorine, bromine or iodine atoms, by C₁₋₅-alkyl, trifluoromethyl, hydroxy, C₁₋₃-alkoxy, carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₄-alkylamino-carbonyl, di-(C₁₋₄-alkyl)-amino-carbonyl, aminosulphonyl, C₁₋₃-alkyl-aminosulphonyl, di-(C₁₋₃-alkyl)-aminosulphonyl, C₁₋₃-alkyl-sulphonylamino, nitro or cyano groups, wherein the substituents may be identical or different, or two adjacent hydrogen atoms of the phenyl groups may be replaced by a methylenedioxy group,

20

and

25

R₅ denotes a hydrogen atom or a C₁₋₃-alkyl group,

wherein by an aryl group is meant a phenyl or naphthyl group optionally mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a cyano, trifluoromethyl, nitro, carboxy, aminocarbonyl, C₁₋₃-alkyl or C₁₋₃-alkoxy group and

30

by a heteroaryl group is meant a monocyclic 5- or 6-membered heteroaryl group optionally substituted by a C₁₋₃-alkyl group in the carbon skeleton, wherein

the 6-membered heteroaryl group contains one, two or three nitrogen atoms and

5

the 5-membered heteroaryl group contains an imino group optionally substituted by a C₁₋₃-alkyl or phenyl-C₁₋₃-alkyl group, an oxygen or sulphur atom or

an imino group optionally substituted by a C₁₋₃-alkyl or phenyl-C₁₋₃-alkyl group or an oxygen or sulphur atom and additionally a nitrogen atom or

10

an imino group optionally substituted by a C₁₋₃-alkyl or phenyl-C₁₋₃-alkyl group and two nitrogen atoms,

15

and moreover a phenyl ring may be fused to the abovementioned monocyclic heterocyclic groups via two adjacent carbon atoms and the bonding takes place via a nitrogen atom or via a carbon atom of the heterocyclic moiety or a fused phenyl ring,

some or all of the hydrogen atoms in the abovementioned alkyl and alkoxy groups or in the alkyl moieties contained in the above-defined groups of formula I optionally being replaced by fluorine atoms,

20

the saturated alkyl and alkoxy moieties with more than 2 carbon atoms which are present in the groups defined hereinbefore also include the branched isomers thereof, such as for example the isopropyl, tert.butyl, isobutyl group, unless otherwise stated, and

25

additionally the hydrogen atom of any carboxy group present or a hydrogen atom bound to a nitrogen atom, e.g. a hydrogen atom of an amino, alkylamino or imino group or a saturated N-heterocycle such as the piperidinyl group, may each be replaced by a group which can be cleaved *in vivo*.

30

By a group which can be cleaved *in vivo* from an imino or amino group is meant, for example, a hydroxy group, an acyl group such as the benzoyl or pyridinoyl group or a C₁₋₁₆-alkanoyl group such as the formyl, acetyl, propionyl, butanoyl, pentanoyl or hexanoyl group, an allyloxycarbonyl group, a C₁₋₁₆-alkoxycarbonyl group such as the methoxy-carbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, 5 tert.butoxycarbonyl, pentoxycarbonyl, hexyloxycarbonyl, octyloxycarbonyl, nonyloxycarbonyl, decyloxycarbonyl, undecyloxycarbonyl, dodecyloxycarbonyl or hexadecyloxycarbonyl group, a phenyl-C₁₋₆-alkoxycarbonyl group such as the benzyloxycarbonyl, phenylethoxycarbonyl or phenylpropoxycarbonyl group, a 10 C₁₋₃-alkylsulphonyl-C₂₋₄-alkoxycarbonyl, C₁₋₃-alkoxy-C₂₋₄-alkoxy-C₂₋₄-alkoxycarbonyl or R_eCO-O-(R_fCR_g)-O-CO group wherein

R_e denotes a C₁₋₈-alkyl, C₅₋₇-cycloalkyl, phenyl or phenyl- C₁₋₃-alkyl group,

15 R_f denotes a hydrogen atom, a C₁₋₃-alkyl, C₅₋₇-cycloalkyl or phenyl group and

R_g denotes a hydrogen atom, a C₁₋₃-alkyl or R_eCO-O-(R_fCR_g)-O group wherein R_e to R_g are as hereinbefore defined,

20 wherein additionally the amino group may be a phthalimido group, whilst the abovementioned ester groups may also be used as a group which can be converted *in vivo* into a carboxy group.

One sub-group of compounds of general formula I which deserves special mention comprises 25 those wherein

X, R₁ and R₃ to R₅ are as hereinbefore defined and

R₂ denotes a straight-chain or branched C₁₋₆-alkoxy-carbonyl group, a 30 C₄₋₇-cycloalkoxycarbonyl or a aryloxycarbonyl group,

a straight-chain or branched C₁₋₆-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, heteroaryl, carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group,

- 5 a straight-chain or branched C₂₋₆-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a chlorine atom or a hydroxy, C₁₋₃-alkoxy, amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts
10 thereof.

A second sub-group of compounds of general formula I which deserves special mention comprises those wherein

- 15 X, R₁ and R₃ to R₅ are as hereinbefore defined and

R₂ denotes an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C₁₋₃-alkoxy group or, if R₄ does not denote an aminosulphonyl-phenyl or N-(C₁₋₅-alkyl)-
20 C₁₋₃-alkylaminocarbonyl-phenyl group, R₂ may also denote a di-(C₁₋₂-alkyl)-aminocarbonyl group,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts
25 thereof.

A third sub-group of compounds of general formula I which deserves special mention comprises those wherein


- 30 X, R₁ to R₃ and R₅ are as hereinbefore defined and

R₄ denotes an R₇-(C₁₋₄-alkyl)-phenyl group, wherein

R₇ denotes an amino, C₁₋₇-alkylamino, di-(C₁₋₇-alkyl)-amino, phenylamino, N-phenyl-C₁₋₃-alkyl-amino, phenyl-C₁₋₃-alkylamino, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino or di-(phenyl-C₁₋₃-alkyl)-amino group,

5

or a phenyl group substituted by the group of formula



10

wherein R₁₂, p and R₁₃ are as hereinbefore defined,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

15

Preferred compounds of general formula I are those wherein

R₁ and R₃ are as hereinbefore defined and

20 X denotes an oxygen atom,

R₂ denotes a carboxy group, a straight-chain or branched C₁₋₆-alkoxy-carbonyl group, a C₅₋₇-cycloalkoxycarbonyl or a phenoxycarbonyl group,

25 a straight-chain or branched C₁₋₃-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, heteroaryl, carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group,

30 a straight-chain or branched C₂₋₃-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a chlorine atom, by a hydroxy, C₁₋₃-alkoxy, amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group,

an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C₁₋₃-alkoxy group or, if R₄ does not denote an aminosulphonyl-phenyl or N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl-phenyl group, it may also denote a di-(C₁₋₂-alkyl)-aminocarbonyl group,

R₄ denotes a C₃₋₇-cycloalkyl group,


whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group or replaced by an -NH or -N(C₁₋₃-alkyl) group,

or a phenyl group substituted by the group R₆, which may additionally be mono- or disubstituted by fluorine, chlorine or bromine atoms, by C₁₋₃-alkyl, trifluoromethyl, hydroxy, C₁₋₃-alkoxy, carboxy, C₁₋₃-alkoxycarbonyl, amino, acetylamino, aminocarbonyl, C₁₋₃-alkyl-aminocarbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, nitro or cyano groups, wherein the substituents may be identical or different and wherein

R₆ denotes a hydrogen, fluorine, chlorine, bromine or iodine atom,

a cyano, nitro, amino, C₁₋₅-alkyl, C₃₋₇-cycloalkyl, trifluoromethyl, phenyl, tetrazolyl or heteroaryl group,

the group of formula



wherein a hydrogen atom bound to the nitrogen atom may be replaced by a C₁₋₃-alkyl group,

a C₁₋₃-alkoxy group, an amino-C₂₋₃-alkoxy, C₁₋₃-alkylamino-C₂₋₃-alkoxy, di-(C₁₋₃-alkyl)-amino-C₂₋₃-alkoxy, phenyl-C₁₋₃-alkylamino-C₂₋₃-alkoxy, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino-C₂₋₃-alkoxy, pyrrolidino-C₂₋₃-alkoxy, piperidino-C₂₋₃-alkoxy or C₁₋₃-alkylmercapto group,

5

a carboxy, C₁₋₄-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylamino-carbonyl, phenyl-C₁₋₃-alkylamino-carbonyl or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino-carbonyl group,

10 a C₃₋₇-cycloalkyl-carbonyl group,

wherein the methylene group in the 4 position of the 6- or 7-membered cycloalkyl moiety may be replaced by an -NH or -N(C₁₋₃-alkyl) group,

15 a 4- to 7-membered cycloalkyleneimino group, wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

20 one or two hydrogen atoms may each be replaced by a C₁₋₃-alkyl group and/or

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, phenyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino group or

25

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH or -N(C₁₋₃-alkyl) group,

30 a C₁₋₄-alkyl group terminally substituted by the group R₇, wherein

R₇ denotes a C₅₋₇-cycloalkyl group,

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be replaced by an -NH or -N(C₁₋₃-alkyl) group or

in a 5- to 7-membered cycloalkyl group a -(CH₂)₂ group may be replaced by a -CO-NH group, a -(CH₂)₃ group may be replaced by a -NH-CO-NH- or a -(CH₂)₄ group may be replaced by a -NH-CO-NH-CO group, whilst in each case a hydrogen atom bound to a nitrogen atom may be replaced by a C₁₋₃-alkyl group,

a phenyl or heteroaryl group,

a hydroxy or C₁₋₃-alkoxy group,

an amino, C₁₋₆-alkylamino, di-(C₁₋₆-alkyl)-amino, phenylamino, N-phenyl-C₁₋₃-alkyl-amino, phenyl-C₁₋₃-alkylamino, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino or di-(phenyl-C₁₋₃-alkyl)-amino group,

a ω-hydroxy-C₂₋₃-alkyl-amino, N-(C₁₋₃-alkyl)-ω-hydroxy--C₂₋₃-alkyl-amino, di-(ω-hydroxy-C₂₋₃-alkyl)-amino, di-(ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl)-amino or N-(dioxolan-2-yl)-C₁₋₃-alkyl-amino group,

a C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-amino or

C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-N-(C₁₋₃-alkyl)-amino group,

a C₁₋₃-alkylsulphonylamino, N-(C₁₋₃-alkyl)-C₁₋₃-alkylsulphonylamino, C₁₋₃-alkylsulphonylamino--C₂₋₃-alkyl-amino or C₁₋₃-alkylsulphonylamino-C₂₋₃-alkyl-N-(C₁₋₃-alkyl)-amino group,

a hydroxycarbonyl- C_{1-3} -alkylamino or
 N-(C_{1-3} -alkyl)-hydroxycarbonyl- C_{1-3} -alkyl-amino group

a guanidino group wherein a hydrogen atom may be replaced by a C_{1-3} -alkyl group,

5

a group of formula



10

wherein

R_8 denotes a hydrogen atom or a C_{1-3} -alkyl group,

n denotes one of the numbers 0, 1, 2 or 3 and

15


R_9 denotes an amino, C_{1-3} -alkylamino, di-(C_{1-3} -alkyl)-amino, phenylamino, benzylamino or C_{1-4} -alkoxy group, a 5- to 7-membered cycloalkyleneimino group, wherein the methylene group in position 4 of the piperidino group may be replaced by an oxygen or sulphur atom, by an -NH, -N(C_{1-3} -alkyl), -N(phenyl), -N(C_{1-3} -alkyl-carbonyl) or -N(benzoyl) group, or, if n denotes one of the

20

numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

25



wherein

R_{10} denotes a hydrogen atom, a C_{1-3} -alkyl group, a C_{1-3} -alkylcarbonyl or
 C_{1-3} -alkylsulphonyl group,

30

m denotes one of the numbers 1, 2 or 3,

o denotes the number 1 or, if m is one of the numbers 2 or 3, o may also denote the number 0 and

5

R₁₁ denotes an amino, C₁₋₃-alkylamino, di-(C₁₋₃-alkyl)-amino, C₁₋₄-alkoxy or C₁₋₃-alkoxy-C₁₋₃-alkoxy group or a 5- to 7-membered cycloalkyleneimino group, wherein the methylene group in position 4 of the piperidino group may be replaced by an oxygen or sulphur atom, by an -NH, -N(C₁₋₃-alkyl), -N(phenyl),
10 -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group,

a C₄₋₇-cycloalkylamino or C₄₋₇-cycloalkenylamino group wherein position 1 of the ring is not involved in the double bond,

15

a 4- to 7-membered cycloalkyleneimino group, wherein

the cycloalkylene moiety may be fused to a phenyl group or

one or two hydrogen atoms may each be replaced by a C₁₋₃-alkyl group and/or

20

the methylene group in position 3 of the pyrrolidino group may be substituted by a hydroxy or C₁₋₃-alkoxy group,

25

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a hydroxy, hydroxy-C₁₋₃-alkyl, C₁₋₃-alkoxy, carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylamino-carbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, phenyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl), -N(phenyl-C₁₋₃-alkyl), -N(C₁₋₃-alkyl-carbonyl), -N(C₁₋₄-alkoxy-carbonyl), -N(benzoyl) or -N(phenyl-C₁₋₃-alkyl-carbonyl) group,

5 wherein a methylene group linked to an imino-nitrogen atom of the cycloalkyleneimino group may be replaced by a carbonyl or sulphonyl group or in a 5- to 6-membered monocyclic cycloalkyleneimino group or a cycloalkyleneimino group fused to a phenyl group the two methylene groups linked to the imino-nitrogen atom may each be replaced by a
10 carbonyl group,

or R₆ denotes a C₁₋₄-alkyl group which is terminally substituted by a carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl or
15 di-(C₁₋₃-alkyl)-aminocarbonyl group or by a 4- to 7-membered cycloalkyleneiminocarbonyl group,

a group of formula

20
$$-N(R_{12})-CO-(CH_2)_p-R_{13} \quad (IV),$$


wherein

R₁₂ denotes a hydrogen atom, a C₁₋₃-alkyl, C₅₋₇-cycloalkyl, phenyl-C₁₋₃-alkyl or
25 heteroaryl-C₁₋₃-alkyl group and

p denotes one of the numbers 0, 1, 2 or 3 and

R₁₃ assumes the meanings of the abovementioned group R₇, or, if p denotes one of
30 the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula



5 wherein


R_{14} denotes a hydrogen atom, a C_{1-4} -alkyl group, a C_{1-3} -alkylcarbonyl, phenylcarbonyl, phenyl- C_{1-3} -alkylcarbonyl, heteroarylcarbonyl, heteroaryl- C_{1-3} -alkylcarbonyl, C_{1-4} -alkylsulphonyl, phenylsulphonyl, phenyl- C_{1-3} -alkylsulphonyl- heteroarylsulphonyl or heteroaryl- C_{1-3} -alkyl-sulphonyl group,

q denotes one of the numbers 1, 2, 3 or 4,

15 r denotes the number 1 or, if q is one of the numbers 2, 3 or 4, it may also denote the number 0 and

R_{15} assumes the meanings of the abovementioned group R_7 ,

20 a group of formula



wherein

25 R_{16} denotes a hydrogen atom or a C_{1-4} -alkyl group optionally terminally substituted by a cyano, trifluoromethyl-carbonylamino or N-(C_{1-3} -alkyl)-trifluoromethyl-carbonyl-amino group and

R_{17} denotes a C_{1-3} -alkyl group,

30

an amino group substituted by a di-(C₁₋₃-alkyl)-amino-C₁₋₃-alkyl-carbonyl or di-(C₁₋₃-alkyl)-amino-C₁₋₃-alkyl-sulphonyl group and a di-(C₁₋₃-alkyl)-aminocarbonyl-C₁₋₃-alkyl group,

5 wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R₆ may be mono- or disubstituted by fluorine, chlorine or bromine atoms, by C₁₋₃-alkyl, trifluoromethyl, hydroxy, C₁₋₃-alkoxy, carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkyl-aminocarbonyl, aminosulphonyl, C₁₋₃-alkyl-aminosulphonyl, nitro or cyano groups, wherein the
10 substituents may be identical or different, or two adjacent hydrogen atoms of the phenyl groups may be replaced by a methylenedioxy group, and

R₅ denotes a hydrogen atom or a C₁₋₃-alkyl group,

15 whilst by a heteroaryl group as mentioned above is meant a pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, furyl, thienyl, oxazolyl, thiazolyl, pyrazolyl, imidazolyl or triazolyl group optionally substituted in the carbon skeleton by a C₁₋₃-alkyl group wherein a hydrogen atom bound to a nitrogen atom may be replaced by a C₁₋₃-alkyl or phenyl-C₁₋₃-alkyl group and wherein the 5-membered heteroaryl groups containing at least
20 one imino group are bound via a carbon or nitrogen atom,

a hydrogen atom bound to a nitrogen atom in the abovementioned groups may be replaced by a group which can be cleaved *in vivo*, particularly by an acetyl or tert.butoxycarbonyl group,

25 the carboxy groups contained in the abovementioned groups may each be substituted by a group which can be cleaved *in vivo* and may occur, for example, in the form of the tert.butoxycarbonyl group,

some or all of the hydrogen atoms in the abovementioned alkyl and alkoxy groups or in the alkyl moieties contained in the above-defined groups of formula I optionally being replaced by fluorine atoms and

- 5 the saturated alkyl and alkoxy moieties contained in the abovementioned groups, which contain more than 2 carbon atoms, may be straight-chain or branched, unless otherwise stated,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts
10 thereof.

One subgroup of preferred compounds of general formula I deserving special mention comprises those wherein

- 15 X, R₁ and R₃ to R₅ are as hereinbefore defined and

R₂ denotes a straight-chain or branched C₁₋₆-alkoxy-carbonyl group, a C₅₋₇-cycloalkoxycarbonyl or a phenoxycarbonyl group,

- 20 a straight-chain or branched C₁₋₃-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl- carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group,

- a straight-chain or branched C₂₋₃-alkoxy-carbonyl group, which is terminally substituted in
25 the alkyl moiety by a hydroxy, C₁₋₃-alkoxy, amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

30

A second sub-group of preferred compounds of general formula I deserving special mention comprises those wherein

X, R₁ and R₃ to R₅ are as hereinbefore defined and

5

R₂ denotes an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C₁₋₃-alkoxy group or, if R₄ does not denote an aminosulphonyl-phenyl or N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl-phenyl group, R₂ may also denote a di-(C₁₋₂-alkyl)-aminocarbonyl group,

10

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

15 A third sub-group of preferred compounds of general formula I deserving special mention comprises those wherein


X, R₁ to R₃ and R₅ are as hereinbefore defined and

20 R₄ denotes an R₇-(n-C₁₋₄-alkyl)-phenyl group, wherein

R₇ denotes an amino, C₁₋₆-alkylamino, di-(C₁₋₆-alkyl)-amino, phenylamino, N-phenyl-C₁₋₃-alkyl-amino, phenyl-C₁₋₃-alkylamino, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino or di-(phenyl-C₁₋₃-alkyl)-amino group,

25

or a phenyl group substituted by the group of formula



30 wherein R₁₂, p and R₁₃ are as hereinbefore defined,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

5 Particularly preferred compounds of general formula I are those wherein

X denotes an oxygen atom,

R₁ denotes a hydrogen atom,

10

R₂ denotes a carboxy group, a straight-chain or branched C₁₋₄-alkoxycarbonyl group or a phenoxycarbonyl group,

15

a straight-chain or branched C₁₋₃-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group,

20

a straight-chain or branched C₂₋₃-alkoxy-carbonyl group which is terminally substituted in the alkyl moiety by a hydroxy, C₁₋₃-alkoxy, amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group,

25

an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C₁₋₃-alkoxy group or, if R₄ does not denote an aminosulphonyl-phenyl or N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl-phenyl group, it may also denote a di-(C₁₋₂-alkyl)-aminocarbonyl group,

30

R₃ denotes a C₁₋₄-alkyl group or a phenyl group which may be substituted by a fluorine, chlorine or bromine atom, by a trifluoromethyl, C₁₋₃-alkyl, hydroxy or C₁₋₃-alkoxy group,

R₄ denotes a C₅₋₆-cycloalkyl group,

wherein the methylene group in position 4 of the cyclohexyl group may be substituted by an amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group or replaced by an -NH or -N(C₁₋₃-alkyl) group,

5


a phenyl group, a phenyl group disubstituted by C₁₋₃-alkyl, C₁₋₃-alkoxy or nitro groups, wherein the substituents may be identical or different, or

a phenyl group substituted by the group R₆, which may additionally be substituted by a
10 fluorine, chlorine or bromine atom or by an amino or nitro group, wherein R₆ denotes a fluorine, chlorine or bromine atom,

a C₁₋₃-alkyl, C₁₋₃-alkoxy, nitro, amino or C₅₋₆-cycloalkyl group,

15 a pyrrolyl, pyrazolyl, imidazolyl, triazolyl or tetrazolyl group bound via a carbon atom, wherein the abovementioned heteroaromatic groups in the carbon skeleton may be substituted by a C₁₋₃-alkyl group or a hydrogen atom bound to a nitrogen atom may be replaced by a C₁₋₃-alkyl or phenyl-C₁₋₃-alkyl group,

20 the group of formula



a carboxy, C₁₋₄-alkoxycarbonyl, phenyl-C₁₋₃-alkylamino-carbonyl or C₅₋₇-cycloalkyl-carbonyl group,

25

a 5 or 6-membered cycloalkyleneimino group, wherein

the methylene group in position 4 of the piperidino group may be replaced by an oxygen or sulphur atom, by an -NH or -N(C₁₋₃-alkyl) group,

an unbranched C₁₋₃-alkyl group terminally substituted by the group R₇, wherein

5

R₇ denotes a C₅₋₇-cycloalkyl group,

10

wherein in a 5 or 6-membered cycloalkyl group a -(CH₂)₂ group may be replaced by a -CO-NH group, a -(CH₂)₃ group may be replaced by an -NH-CO-NH- or a -(CH₂)₄ group may be replaced by an -NH-CO-NH-CO group, whilst in each case a hydrogen atom bound to a nitrogen atom may be replaced by a C₁₋₃-alkyl group,

15

a phenyl or pyridinyl group or a pyrrolyl, pyrazolyl, imidazolyl or triazolyl group bound via a carbon or nitrogen atom, wherein the abovementioned heteroaromatic groups in the carbon skeleton may be substituted by a C₁₋₃-alkyl group or a hydrogen atom bound to a nitrogen atom may be replaced by a C₁₋₃-alkyl group,

20

a hydroxy or C₁₋₃-alkoxy group,

25

an amino, C₁₋₆-alkylamino, di-(C₁₋₆-alkyl)-amino, phenylamino, N-phenyl-C₁₋₃-alkylamino, phenyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino group,

a ω-hydroxy-C₂₋₃-alkyl-amino, N-(C₁₋₃-alkyl)-ω-hydroxy-C₂₋₃-alkylamino, di-(ω-hydroxy-C₂₋₃-alkyl)-amino or di-(ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl)-amino group,

30

a C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-amino or C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-N-(C₁₋₃-alkyl)-amino group,

a C₁₋₃-alkylsulphonylamino, N-(C₁₋₃-alkyl)-C₁₋₃-alkylsulphonylamino,
C₁₋₃-alkylsulphonylamino--C₂₋₃-alkylamino or C₁₋₃-alkylsulphonylamino-
-C₂₋₃-alkyl-N-(C₁₋₃-alkyl)-amino group,

5 a hydroxycarbonyl-C₁₋₃-alkylamino or
N-(C₁₋₃-alkyl)-hydroxycarbonyl-C₁₋₃-alkyl-amino group,

a guanidino group wherein a hydrogen atom may be replaced by a C₁₋₃-alkyl group,

10 a group of formula




wherein

15 R₈ denotes a hydrogen atom or a C₁₋₃-alkyl group,

n denotes one of the numbers 0, 1, 2 or 3 and

20 R₉ denotes an amino, C₁₋₃-alkylamino, di-(C₁₋₃-alkyl)-amino or C₁₋₄-alkoxy
group, a 5- or 6-membered cycloalkyleneimino group, wherein the methylene
group in position 4 of the piperidino group may be replaced by an -NH,
-N(C₁₋₃-alkyl) or -N(C₁₋₃-alkyl-carbonyl) group, or, if n denotes one of the
numbers 1, 2 or 3, R₉ may also denote a hydrogen atom,

25 a group of formula



30 wherein

R_{10} denotes a hydrogen atom or a C_{1-3} -alkyl group,

m denotes one of the numbers 1, 2 or 3,

5 o denotes the number 1 or, if m is one of the numbers 2 or 3, o may also denote the number 0 and

10 R_{11} denotes an amino, C_{1-3} -alkylamino, di- $(C_{1-3}$ -alkyl)-amino, C_{1-4} -alkoxy or methoxy- C_{1-3} -alkoxy group or a 5- or 6-membered cycloalkyleneimino group, wherein the methylene group in position 4 of the piperidino group may be replaced by an -NH, -N(C_{1-3} -alkyl) or -N(C_{1-3} -alkyl-carbonyl) group,

an azetidino, pyrrolidino, piperidino, 2,6-dimethyl-piperidino, 3,5-dimethyl-piperidino or azepino group, wherein

15 the methylene group in position 3 of the pyrrolidino group may be substituted by a hydroxy group,


20 the methylene group in position 4 of the piperidino group may be substituted by a hydroxy, hydroxy- C_{1-3} -alkyl or C_{1-3} -alkoxy group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3} -alkyl), -N(C_{1-3} -alkyl-carbonyl), -N(benzoyl) or -N(phenyl- C_{1-3} -alkyl-carbonyl) group,

25 wherein a methylene group linked to an imino-nitrogen atom of the pyrrolidino, piperidino or piperazino group may be replaced by a carbonyl group,

30 or R_6 denotes a straight-chain C_{1-3} -alkyl group which is terminally substituted by a carboxy or C_{1-3} -alkoxy-carbonyl group,

a group of formula



5

wherein

R_{12} denotes a hydrogen atom, a C_{1-3} -alkyl or phenyl- C_{1-3} -alkyl group,

10 p denotes one of the numbers 0, 1 or 2 and

R_{13} denotes an amino, C_{1-4} -alkylamino, di- $(C_{1-4}$ -alkyl)-amino, benzylamino, N- $(C_{1-3}$ -alkyl)-benzylamino, C_{1-3} -alkoxy- C_{1-3} -alkylamino, N- $(C_{1-3}$ -alkyl)- C_{1-3} -alkoxy- C_{1-3} -alkylamino, di-(2-methoxy-ethyl)-amino, di-(ω -hydroxy- C_{2-3} -alkyl)-amino or aminocarbonyl-methyl-N-(methyl)-amino group,


15

a pyrrolyl, pyrazolyl or imidazolyl group bound via a nitrogen atom and optionally substituted by a C_{1-3} -alkyl group,

20

a pyrrolidino, piperidino, morpholino, thiomorpholino or a piperazino group optionally substituted in the 4 position by a C_{1-3} -alkyl, phenyl- C_{1-3} -alkyl, C_{1-3} -alkylcarbonyl or C_{1-4} -alkoxycarbonyl group or, if n denotes the number 1 or 2, it may also denote a hydrogen atom,

25 a group of formula



wherein

30

R₁₄ denotes a hydrogen atom, a C₁₋₄-alkyl, C₁₋₃-alkyl-carbonyl, phenylcarbonyl, phenyl-C₁₋₃-alkylcarbonyl, furyl-carbonyl, pyridinyl-carbonyl, furyl-C₁₋₃-alkyl-carbonyl, pyridinyl-C₁₋₃-alkylcarbonyl, C₁₋₄-alkylsulphonyl, phenylsulphonyl or phenyl-C₁₋₃-alkylsulphonyl group,

5

q denotes one of the numbers 1, 2 or 3,


r denotes the number 1 or, if q is one of the numbers 2 or 3, it may also denote the number 0 and

10

R₁₅ denotes an amino, C₁₋₄-alkylamino, di-(C₁₋₄-alkyl)-amino, phenylamino, N-(C₁₋₄-alkyl)-phenylamino, benzylamino or N-(C₁₋₄-alkyl)-benzylamino group,

or a group of formula

15



wherein

R₁₆ denotes a hydrogen atom or a C₁₋₃-alkyl group optionally terminally substituted by a cyano, trifluoromethyl-carbonylamino or N-(C₁₋₃-alkyl)-trifluoromethyl-carbonyl-amino group and

20

R₁₇ denotes a C₁₋₃-alkyl group,

25

wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R₆ may be substituted by a fluorine, chlorine or bromine atom, by a methyl, trifluoromethyl, methoxy, nitro or cyano group and

R₅ denotes a hydrogen atom,

30

wherein a hydrogen atom bound to a nitrogen atom in the abovementioned groups may be replaced by an acetyl or tert.butoxycarbonyl group,

the carboxy groups contained in the abovementioned groups may also be present in the
5 form of the tert.butoxycarbonyl precursor group and

the saturated alkyl and alkoxy moieties contained in the abovementioned groups, which contain more than 2 carbon atoms, may be straight-chain or branched, unless otherwise stated,

10

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

15

One subgroup of particularly preferred compounds of general formula I deserving special mention comprises those wherein

X, R₁, R₃ and R₅ are as hereinbefore defined,

20

R₂ denotes a straight-chain or branched C₁₋₄-alkoxycarbonyl group or a phenoxycarbonyl group,

25

a straight-chain or branched C₁₋₃-alkoxycarbonyl group, which is terminally substituted in the alkyl moiety by a phenyl- carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group, or


a straight-chain or branched C₂₋₃-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a hydroxy, C₁₋₃-alkoxy, amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group, and

30

R₄ denotes an R₇-(n-C₁₋₃-alkyl)-phenyl group, wherein

R₇ denotes an amino, C₁₋₆-alkylamino, di-(C₁₋₄-alkyl)-amino,
 ω-hydroxy-C₂₋₃-alkyl-amino, N-(C₁₋₃-alkyl)-ω-hydroxy-C₂₋₃-alkyl-amino,
 di-(ω-hydroxy-C₂₋₃-alkyl)-amino or di-(ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl)-amino group,

5 or a phenyl group substituted by the group of formula



wherein R₁₂, p and R₁₃ are as hereinbefore defined,

10

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

15

A second subgroup of particularly preferred compounds of general formula I deserving special mention comprises those wherein

X, R₁, R₃ and R₅ are as hereinbefore defined,

20


R₂ denotes an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C₁₋₃-alkoxy group or, if R₄ does not denote an aminosulphonyl-phenyl or N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl-phenyl group, R₂ may also denote a di-(C₁₋₂-alkyl)-aminocarbonyl group and

25

R₄ denotes a R₇-(n-C₁₋₃-alkyl)-phenyl group, wherein

30

R₇ denotes an amino, C₁₋₆-alkylamino, di-(C₁₋₄-alkyl)-amino,
 ω-hydroxy-C₂₋₃-alkyl-amino, N-(C₁₋₃-alkyl)-ω-hydroxy-C₂₋₃-alkyl-amino,
 di-(ω-hydroxy-C₂₋₃-alkyl)-amino or di-(ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl)-amino group,
 or a phenyl group substituted by the group of formula



wherein R_{12} , p and R_{13} are as hereinbefore defined,

5

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

Most particularly preferred compounds of general formula I are those wherein

10

X denotes an oxygen atom,

R_1 and R_5 each denote a hydrogen atom,

15

R_2 denotes a methoxycarbonyl, ethoxycarbonyl or aminocarbonyl group,

R_3 denotes a phenyl group and

R_4 denotes a phenyl group monosubstituted by the group R_6 , wherein


20

R_6 denotes an N-methyl-imidazol-2-yl group,

an unbranched C_{1-3} -alkyl group which is terminally substituted by a C_{1-4} -alkylamino, di- $(C_{1-4}$ -alkyl)-amino, piperidino or 2,6-dimethyl-piperidino group,

25

a group of formula



30


wherein

R₁₂ denotes a C₁₋₃-alkyl group,

p denotes one of the numbers 1 or 2 and

5 R₁₃ denotes a di-(C₁₋₃-alkyl)-amino group,

or a group of formula



10

wherein

R₁₄ denotes a C₁₋₃-alkyl-carbonyl or C₁₋₃-alkylsulphonyl group,

15 q denotes one of the numbers 1, 2 or 3,

r denotes the number 1 or, if q is one of the numbers 2 or 3, r may also denote the number 0 and

20 R₁₅ denotes a di-(C₁₋₃-alkyl)-amino group,

wherein the saturated alkyl moieties contained in the abovementioned groups which contain more than 2 carbon atoms may be straight-chain or branched, unless otherwise stated,

25

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.


A subgroup of most particularly preferred compounds of general formula I deserving special
30 mention comprises those wherein

X, R₁, R₃ and R₅ are as hereinbefore defined,

R₂ denotes a methoxycarbonyl or ethoxycarbonyl group and

5 R₄ denotes a di-(C₁₋₃-alkyl)-amino-C₁₋₃-alkylphenyl group or

a phenyl group substituted by the group of formula



10

wherein R₁₂, p and R₁₃ are as hereinbefore defined,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

15

The following are mentioned as examples of particularly preferred compounds:

(a) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

20

(b) 3-Z-[(1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone,

(c) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

25

(d) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

30 (e) 3-Z-[1-(4-((2,6-dimethyl-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(f) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

5 (g) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(h) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

10

(i) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(j) 3-Z-[1-(4-(N-acetyl-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

15

(k) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

20 (l) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(m) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

25

(n) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(o) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

30

(p) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(q) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(r) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone and

(s) 3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

the tautomers, the mixtures and the salts thereof.

Another subgroup of compounds of general formula I comprises those wherein X denotes an oxygen or sulphur atom,

R₁ denotes a hydrogen atom or a prodrug group such as a C₁₋₄-alkoxycarbonyl or C₂₋₄-alkanoyl group,

R₂ denotes a carboxy group, a straight-chain or branched C₁₋₆-alkoxycarbonyl group, a C₅₋₇-cycloalkoxycarbonyl or phenyl-C₁₋₃-alkoxycarbonyl group, an aminocarbonyl or C₁₋₂-alkylaminocarbonyl group or, if R₄ does not denote an aminosulphonyl-phenyl or N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl-phenyl group, a di-(C₁₋₂-alkyl)-aminocarbonyl group,

R₃ denotes a hydrogen atom, a C₁₋₆-alkyl, C₃₋₇-cycloalkyl, trifluoromethyl or heteroaryl group,

a phenyl or naphthyl group, a phenyl or naphthyl group mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a trifluoromethyl, C₁₋₃-alkyl or C₁₋₃-alkoxy

group, whilst in the event of disubstitution the substituents may be identical or different and wherein the abovementioned unsubstituted as well as the mono- and disubstituted phenyl and naphthyl groups may additionally be substituted

5 by a hydroxy, hydroxy-C₁₋₃-alkyl or C₁₋₃-alkoxy-C₁₋₃-alkyl group,

by a cyano, carboxy, carboxy-C₁₋₃-alkyl, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylamino-carbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group,

10 by a nitro group,

by an amino, C₁₋₃-alkylamino, di-(C₁₋₃-alkyl)-amino or amino-C₁₋₃-alkyl group,

15 by a C₁₋₃-alkylcarbonylamino, N-(C₁₋₃-alkyl)-C₁₋₃-alkyl-carbonylamino, C₁₋₃-alkylcarbonylamino-C₁₋₃-alkyl, N-(C₁₋₃-alkyl)-C₁₋₃-alkylcarbonylamino-C₁₋₃-alkyl, C₁₋₃-alkyl-sulphonylamino, C₁₋₃-alkylsulphonylamino-C₁₋₃-alkyl, N-(C₁₋₃-alkyl)-C₁₋₃-alkylsulphonylamino-C₁₋₃-alkyl or aryl-C₁₋₃-alkylsulphonylamino group,

20 by a cycloalkylamino, cycloalkyleneimino, cycloalkyleneiminocarbonyl, cycloalkyleneimino-C₁₋₃-alkyl, cycloalkyleneiminocarbonyl-C₁₋₃-alkyl or cycloalkyleneiminosulphonyl-C₁₋₃-alkyl group having 4 to 7 ring members in each case, whilst in each case the methylene group in position 4 of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a
25 sulphinyl, sulphonyl, -NH or -N(C₁₋₃-alkyl) group,

or by a heteroaryl or heteroaryl-C₁₋₃-alkyl group,

R₄ denotes a C₃₋₇-cycloalkyl group,

30

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group or replaced by an -NH or -N(C₁₋₃-alkyl) group,

- 5 or a phenyl group substituted by the group R₆, which may additionally be substituted by a fluorine, chlorine, bromine or iodine atom, by a C₁₋₅-alkyl, trifluoromethyl, C₁₋₃-alkoxy, carboxy, C₁₋₃-alkoxycarbonyl, aminosulphonyl, nitro or cyano group, wherein

R₆ denotes a hydrogen, fluorine, chlorine, bromine or iodine atom,

10

a cyano, nitro, C₁₋₅-alkyl, C₃₋₇-cycloalkyl, trifluoromethyl, phenyl, tetrazolyl or heteroaryl group,

15

a C₁₋₃-alkoxy group optionally substituted by 1 to 3 fluorine atoms, a C₁₋₃-alkoxy-C₁₋₃-alkoxy, phenyl-C₁₋₃-alkoxy, amino-C₂₋₃-alkoxy, C₁₋₃-alkylamino-C₂₋₃-alkoxy, di-(C₁₋₃-alkyl)-amino-C₂₋₃-alkoxy, phenyl-C₁₋₃-alkylamino-C₂₋₃-alkoxy, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino-C₂₋₃-alkoxy, C₅₋₇-cycloalkyleneimino-C₂₋₃-alkoxy or C₁₋₃-alkylmercapto group,

20

a carboxy, C₁₋₄-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylamino-carbonyl, N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl, phenyl-C₁₋₃-alkylamino-carbonyl, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino-carbonyl, piperazinocarbonyl or N-(C₁₋₃-alkyl)-piperazinocarbonyl group,

25

a C₁₋₃-alkylaminocarbonyl or N-(C₁₋₅-alkyl)-C₁₋₃-alkylaminocarbonyl group wherein an alkyl moiety is substituted by a carboxy or C₁₋₃-alkoxycarbonyl group or is substituted in the 2 or 3 position by a di-(C₁₋₃-alkyl)-amino, piperazino, N-(C₁₋₃-alkyl)-piperazino or a 4- to 7-membered cycloalkyleneimino group,

30

a 4- to 7-membered cycloalkyleneimino group, wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

the cycloalkylene moiety may be fused to a phenyl ring or

5

one or two hydrogen atoms may each be replaced by a C₁₋₃-alkyl group and/or

10

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, phenyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino group or

15

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl), -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group,

a C₁₋₄-alkyl group which may be substituted

by a hydroxy or C₁₋₃-alkoxy group,

20

by an amino, C₁₋₇-alkylamino, di-(C₁₋₇-alkyl)-amino, di-N-(C₁₋₃-alkyl)-amino-C₂₋₃-alkylamino, tri-N,N,N'-(C₁₋₃-alkyl)-amino-C₂₋₃-alkylamino, phenylamino, N-phenyl-C₁₋₃-alkyl-amino, phenyl-C₁₋₃-alkylamino, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino or di-(phenyl-C₁₋₃-alkyl)-amino group,

25

by a C₁₋₃-alkylcarbonylamino, N-(C₁₋₃-alkyl)-C₁₋₃-alkylcarbonylamino, C₁₋₃-alkoxycarbonyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-C₁₋₃-alkoxycarbonyl-C₁₋₃-alkylamino group,

30

by a C₄₋₇-cycloalkylamino, C₄₋₇-cycloalkyl-C₁₋₃-alkylamino or C₄₋₇-cycloalkenylamino group wherein position 1 of the ring is not involved in the double bond and wherein the abovementioned groups may each additionally be

substituted at the amino-nitrogen atom by a C₁₋₃-alkyl group wherein some or all of the hydrogen atoms are replaced by fluorine atoms, by a C₅₋₇-cycloalkyl, C₂₋₄-alkenyl or C₁₋₄-alkyl group,

5 by a 4- to 7-membered cycloalkyleneimino group, wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

10 the cycloalkylene moiety may be fused to a phenyl group or to an oxazolo, imidazolo, thiazolo, pyridino, pyrazino or pyrimidino group optionally substituted by a fluorine, chlorine, bromine or iodine atom, by a nitro, C₁₋₃-alkyl, C₁₋₃-alkoxy or amino group or

15 one or two hydrogen atoms may each be replaced by a C₁₋₃-alkyl, C₅₋₇-cycloalkyl or phenyl group and/or

20 in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a hydroxy, carboxy, C₁₋₄-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, phenyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino group or

25 may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl), -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group,

by a carboxy, C₁₋₃-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group or

30 by a 4- to 7-membered cycloalkyleneiminocarbonyl group,

an amino, pyrrolidino, piperidino, morpholino, benzoylamino or N-(C₁₋₃-alkyl)-benzoylamino group,

an N-(C₁₋₃-alkyl)-C₂₋₄-alkanoylamino group which is additionally substituted in the alkyl
5 moiety by a carboxy or C₁₋₃-alkoxycarbonyl group,

a group of formula



10

wherein

R₈ denotes a hydrogen atom or a C₁₋₃-alkyl group,

15


n denotes one of the numbers 0, 1, 2 or 3 and

20

R₉ denotes an amino, C₁₋₄-alkylamino, phenylamino, N-(C₁₋₄-alkyl)-phenylamino, benzylamino, N-(C₁₋₄-alkyl)-benzylamino or di-(C₁₋₄-alkyl)-amino group, a 4- to 7-membered cycloalkyleneimino group, whilst in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl), -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group, or, if n denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

25

a group of formula



wherein

30

R₁₀ denotes a hydrogen atom, a C₁₋₃-alkyl group, a C₁₋₃-alkylcarbonyl, arylcarbonyl, phenyl-C₁₋₃-alkylcarbonyl, C₁₋₃-alkylsulphonyl, arylsulphonyl or phenyl-C₁₋₃-alkylsulphonyl group,

5 m denotes one of the numbers 1, 2, 3 or 4,

o denotes one of the numbers 0 or 1 and

10 R₁₁ denotes an amino, C₁₋₄-alkylamino, phenylamino, N-(C₁₋₄-alkyl)-phenylamino, benzylamino, N-(C₁₋₄-alkyl)-benzylamino or di-(C₁₋₄-alkyl)-amino group, a 4- to 7-membered cycloalkyleneimino group, wherein the cycloalkylene moiety may be fused to a phenyl ring or in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl),
15 -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group, a C₁₋₃-alkoxy group or a di-(C₁₋₄-alkyl)-amino-C₁₋₃-alkylamino group optionally substituted in the 1 position by a C₁₋₃-alkyl group,

20 or an N-(C₁₋₃-alkyl)-C₁₋₅-alkylsulphonylamino or N-(C₁₋₃-alkyl)-phenylsulphonylamino group wherein the alkyl moiety is additionally substituted by a cyano or carboxy group,

25 wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R₆ may be mono- or disubstituted by fluorine, chlorine, bromine or iodine atoms, by C₁₋₅-alkyl, trifluoromethyl, C₁₋₃-alkoxy, carboxy, C₁₋₃-alk-oxycarbonyl, aminosulphonyl, nitro or cyano groups, wherein the substituents may be identical or different, or two adjacent hydrogen atoms of the phenyl groups may be replaced by a methylenedioxy group,

and

30

R₅ denotes a hydrogen atom or a C₁₋₃-alkyl group,

wherein by an aryl group is meant a phenyl or naphthyl group optionally mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a trifluoromethyl, C₁₋₃-alkyl or C₁₋₃-alkoxy group and

5

by a heteroaryl group is meant a monocyclic 5- or 6-membered heteroaryl group optionally substituted by a C₁₋₃-alkyl group, wherein the 6-membered heteroaryl group contains one, two or three nitrogen atoms and the 5-membered heteroaryl group contains an imino group optionally substituted by a C₁₋₃-alkyl group, an oxygen or sulphur atom or an imino group optionally substituted by a C₁₋₃-alkyl group and an oxygen or sulphur atom or one or two
10 nitrogen atoms, and moreover a phenyl ring may be fused to the abovementioned monocyclic heterocyclic groups via two adjacent carbon atoms,

the saturated alkyl and alkoxy moieties present in the groups defined above which contain
15 more than 2 carbon atoms also include the branched isomers thereof such as, for example, the isopropyl, tert.butyl or isobutyl group, unless otherwise stated, and

additionally any carboxy, amino or imino group present may be substituted by a group which can be cleaved *in vivo*,


20

the isomers and the salts thereof.

According to the invention the new compounds are obtained, for example, by the following methods known in principle from the literature:

25

a. reacting a compound of general formula



wherein

X and R₃ are as hereinbefore defined,

R₂' has the meanings given for R₂ hereinbefore,

- 5 R₁₈ denotes a hydrogen atom or a protecting group for the nitrogen atom of the lactam group, wherein one of the groups R₂' and R₁₈ may also denote a bond to a solid phase optionally formed via a spacer and the other one of the groups R₂' and R₁₈ has the abovementioned meanings, and Z₁ denotes a halogen atom, a hydroxy, alkoxy or aryl-alkoxy group, e.g. a chlorine or bromine atom, a methoxy, ethoxy or benzyloxy group,

10

with an amine of general formula



- 15 wherein

R₄ and R₅ are as hereinbefore defined,

and if necessary subsequently cleaving any protecting group used for the nitrogen atom of the lactam group or cleaving from a solid phase.

- 20 The protecting group for the nitrogen atom of the lactam group may be, for example, an acetyl, benzoyl, ethoxycarbonyl, tert.butyloxycarbonyl or benzyloxycarbonyl group and

the solid phase may be a resin such as a 4-(2',4'-dimethoxyphenylaminomethyl)-phenoxy resin, the bond preferably being formed via the amino group, or a p-benzyloxybenzyl alcohol resin, wherein the bond is conveniently formed via an intermediate member such as a 2,5-dimethoxy-4-hydroxy-benzyl derivative.

5

The reaction is conveniently carried out in a solvent such as dimethylformamide, toluene, acetonitrile, tetrahydrofuran, dimethylsulphoxide, methylene chloride or mixtures thereof, optionally in the presence of an inert base such as triethylamine, N-ethyl-diisopropylamine or sodium hydrogen carbonate at temperatures between 20 and 175°C, whilst any protecting group used can be cleaved at the same time by transamidation.

10

If Z_1 in a compound of general formula VII denotes a halogen atom, the reaction is preferably carried out in the presence of an inert base at temperatures of between 20 and 120°C.

15

If Z_1 in a compound of general formula VII denotes a hydroxy, alkoxy or arylalkoxy group, the reaction is preferably carried out at temperatures between 20 and 200°C.

20

If a protecting group used subsequently has to be cleaved, this is conveniently done either hydrolytically in an aqueous or alcoholic solvent, e.g. in methanol/water, ethanol/water, isopropanol/water, tetrahydrofuran/water, dioxan/water, dimethylformamide/water, methanol or ethanol in the presence of an alkali metal base such as lithium hydroxide, sodium hydroxide or potassium hydroxide at temperatures between 0 and 100°C, preferably at temperatures between 10 and 50°C,

25


or advantageously by transamidation with an organic base such as ammonia, butylamine, dimethylamine or piperidine in a solvent such as methanol, ethanol, dimethylformamide and the mixtures thereof or in an excess of the amine used, at temperatures between 0 and 100°C, preferably at temperatures between 10 and 50°C.

30

Cleaving from any solid phase used is preferably carried out using trifluoroacetic acid and water at temperatures between 0 and 35°C, preferably at ambient temperature.

- b. In order to prepare a compound of general formula I wherein R₂ has the meanings given
5 hereinbefore, with the exception of the carboxy group:

reacting a compound of general formula




10

wherein

R₁ and R₃ to R₅ are as hereinbefore defined, or the reactive derivatives thereof, with a compound of general formula

15



wherein

R₁₉ denotes a C₁₋₆-alkanol, a C₄₋₇-cycloalkanol or an aromatic alcohol,

20

a C₁₋₆-alkanol which is terminally substituted in the alkyl moiety by a phenyl, heteroaryl, carboxy, C₁₋₃-alkoxy-carbonyl, aminocarbonyl, C₁₋₃-alkylamino-carbonyl or di-(C₁₋₃-alkyl)-aminocarbonyl group,

25

a C₂₋₆-alkanol which is terminally substituted in the alkyl moiety by a chlorine atom or a hydroxy, C₁₋₃-alkoxy, amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)-amino group,

an amino or methylamino group, an ethylamino group optionally substituted in the 2 position of the ethyl group by a hydroxy or C₁₋₃-alkoxy group or a di-(C₁₋₂-alkyl)-amino group.

5

The esterification or amidation is preferably carried out in a solvent such as methylene chloride, diethylether, tetrahydrofuran, toluene, dioxan, acetonitrile, dimethylsulphoxide or dimethylformamide, optionally in the presence of an inorganic or a tertiary organic base, preferably at temperatures between 20°C and the boiling temperature of the solvent used.

- 10 The reaction with a corresponding acid is preferably carried out in the presence of a dehydrating agent, e.g. in the presence of isobutyl chloroformate, tetraethyl orthocarbonate, trimethyl orthoacetate, 2,2-dimethoxypropane, tetramethoxysilane, thionylchloride, trimethylchlorosilane, phosphorus trichloride, phosphorus pentoxide, N,N'-dicyclohexylcarbodiimide, N,N'-dicyclohexyl-carbodiimide/N-hydroxysuccinimide,
- 15 N,N'-dicyclohexyl-carbodiimide/1-hydroxy-benzotriazole, 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium-tetrafluoroborate, 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium-tetrafluoroborate/1-hydroxy-benzotriazole, N,N'-carbonyldiimidazole or triphenylphosphine/carbon tetrachloride, and optionally with the addition of a base such as pyridine, 4-dimethylaminopyridine, N-methyl-morpholine or triethylamine,
- 20 conveniently at temperatures between 0 and 150°C, preferably at temperatures between 0 and 100°C, and the acylation with a corresponding reactive compound such as an anhydride, ester, imidazolide or halide thereof, is optionally carried out in the presence of a tertiary organic base such as triethylamine, N-ethyl-diisopropylamine or N-methyl-morpholine at temperatures between 0 and 150°C, preferably at temperatures between 50
- 25 and 100°C.

c. In order to prepare a compound of general formula I, wherein R₄ denotes a C₁₋₄-alkyl group substituted by the group R₇, wherein


R₇ denotes an amino, C₁₋₇-alkylamino, di-(C₁₋₇-alkyl)-amino, phenylamino, N-phenyl-C₁₋₃-alkyl-amino, phenyl-C₁₋₃-alkylamino, N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino or di-(phenyl-C₁₋₃-alkyl)-amino group,

5 a ω-hydroxy-C₂₋₃-alkyl-amino, N-(C₁₋₃-alkyl)-ω-hydroxy-C₂₋₃-alkyl-amino, di-(ω-hydroxy-C₂₋₃-alkyl)-amino, di-(ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl)-amino or N-(dioxolan-2-yl)-C₁₋₃-alkyl-amino group,

10 a C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-amino or C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-N-(C₁₋₃-alkyl)-amino group,

a C₁₋₃-alkylsulphonylamino, N-(C₁₋₃-alkyl)-C₁₋₃-alkylsulphonylamino, C₁₋₃-alkylsulphonylamino-C₂₋₃-alkyl-amino or C₁₋₃-alkylsulphonylamino-C₂₋₃-alkyl-N-(C₁₋₃-alkyl)-amino group,

15 a group of formula



20 wherein

R₁₀ denotes a hydrogen atom, a C₁₋₃-alkyl group, a C₁₋₃-alkylcarbonyl, arylcarbonyl, phenyl-C₁₋₃-alkylcarbonyl, C₁₋₃-alkylsulphonyl, arylsulphonyl or phenyl-C₁₋₃-alkylsulphonyl group,

25 m denotes one of the numbers 1, 2, 3 or 4,

o denotes the number 1 and

30 R₁₁ denotes an amino, C₁₋₄-alkylamino, di-(C₁₋₄-alkyl)-amino, phenylamino, N-(C₁₋₄-alkyl)-phenylamino, benzylamino, N-(C₁₋₄-alkyl)-benzylamino,

C₁₋₄-alkoxy or C₁₋₃-alkoxy-C₁₋₃-alkoxy group, a di-(C₁₋₄-alkyl)-amino-C₁₋₃-alkylamino group optionally substituted in the 1 position by a C₁₋₃-alkyl group, or a 4- to 7-membered cycloalkyleneimino group, wherein the cycloalkylene moiety may be fused to a phenyl ring or in each case the methylene group in the
 5 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl), -N(phenyl), -N(C₁₋₃-alkyl-carbonyl) or -N(benzoyl) group,

a C₄₋₇-cycloalkylamino, C₄₋₇-cycloalkyl-C₁₋₃-alkylamino or C₄₋₇-cycloalkenylamino
 10 group wherein position 1 of the ring is not involved in the double bond and wherein the abovementioned groups may each additionally be substituted at the amino-nitrogen atom by a C₅₋₇-cycloalkyl, C₂₋₄-alkenyl or C₁₋₄-alkyl group,

or a 4- to 7-membered cycloalkyleneimino group, wherein

15 the cycloalkylene moiety may be fused to a phenyl group or to an oxazolo, imidazolo, thiazolo, pyridino, pyrazino or pyrimidino group optionally substituted by a fluorine, chlorine, bromine or iodine atom, by a nitro, C₁₋₃-alkyl, C₁₋₃-alkoxy or amino group, and/or

20 one or two hydrogen atoms may each be replaced by a C₁₋₃-alkyl, C₅₋₇-cycloalkyl or phenyl group and/or

25 the methylene group in the 3 position of a 5-membered cycloalkyleneimino group may be substituted by a hydroxy, hydroxy-C₁₋₃-alkyl, C₁₋₃-alkoxy or C₁₋₃-alkoxy-C₁₋₃-alkyl group,


30 in each case the methylene group in the 3 or 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a hydroxy, hydroxy-C₁₋₃-alkyl, C₁₋₃-alkoxy, C₁₋₃-alkoxy-C₁₋₃-alkyl, C₁₋₄-alkoxycarbonyl, aminocarbonyl,

C₁₋₃-alkylaminocarbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, phenyl-C₁₋₃-alkylamino or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C₁₋₃-alkyl-), -N(phenyl), -N(phenyl-C₁₋₃-alkyl-), -N(C₁₋₃-alkyl-carbonyl-), -N(C₁₋₄-alkoxy-carbonyl-), -N(benzoyl-) or -N(phenyl-C₁₋₃-alkyl-carbonyl-) group,

wherein a methylene group linked to an imino-nitrogen atom of the cycloalkyleneimino group may be replaced by a carbonyl or sulphonyl group or in a 5- to 7-membered monocyclic cycloalkyleneimino group or a cycloalkyleneimino group fused to a phenyl group the two methylene groups linked to the imino-nitrogen atom may each be replaced by a carbonyl group:

reacting a compound of general formula



wherein

R₃, R₅ and X are as hereinbefore defined,


R₂' has the meanings given for R₂ hereinbefore,

R₁₈ denotes a hydrogen atom or a protecting group for the nitrogen atom of the lactam group, wherein one of the groups R₂' and R₁₈ may also denote a bond to a solid phase

optionally formed via a spacer and the other one of the groups R₂' and R₁₈ has the abovementioned meanings, A denotes a C₁₋₄-alkyl group and Z₂ denotes a leaving group,

for example an alkyl or arylsulphonyloxy group such as the methylsulphonyloxy, ethylsulphonyloxy, p-toluenesulphonyloxy or trifluoromethanesulphonyloxy group, with an amine of general formula

5



wherein

R_7 has the meanings given for R_7 hereinbefore, and subsequently, if necessary, cleaving
10 any protecting group used for the nitrogen atom of the lactam group, or cleaving from a solid phase.

The reaction is conveniently carried out in a solvent such as methylene chloride, tetrahydrofuran, 1,4-dioxan, toluene, acetonitrile, dimethylsulphoxide, dimethylformamide,
15 dimethylacetamide, N-methylpyrrolidone or the mixtures thereof, optionally with the addition of water as a co-solvent and/or with the addition of an inert auxiliary base, e.g. sodium hydrogen carbonate, pyridine, 2,4,6-trimethylpyridine, quinoline, triethylamine, N-ethyl-diisopropylamine, N-ethyl-dicyclohexylamine, 1,4-diazabicyclo[2,2,2]octane or 1,8-diazabicyclo[5,4,0]undec-7-ene, at temperatures between -50°C and $+100^\circ\text{C}$, preferably
20 between -10°C and $+50^\circ\text{C}$, while any protecting group used may be cleaved at the same time by transamidation.

If any protecting group used for the nitrogen atom of the lactam group has to be removed or if the compound has to be cleaved from a solid phase this is carried out as described
25 under method (a) above.

If according to the invention a compound of general formula I is obtained which contains an alkoxy carbonyl group, this may be converted by hydrolysis into a corresponding carboxy compound, or

30

if a compound of general formula I is obtained which contains an amino or alkylamino group, this may be converted by reductive alkylation into a corresponding alkylamino or dialkylamino compound, or

- 5 if a compound of general formula I is obtained which contains an amino or alkylamino group, this may be converted by acylation or sulphonation into a corresponding acyl or sulphonyl compound, or

- 10 if a compound of general formula I is obtained which contains a carboxy group, this may be converted by esterification or amidation into a corresponding ester or aminocarbonyl compound, or

- 15 if a compound of general formula I is obtained which contains a cycloalkyleneimino group wherein a methylene group is replaced by a sulphur atom, this may be converted by oxidation into a corresponding sulphinyl or sulphonyl compound, or

if a compound of general formula I is obtained which contains a nitro group, this may be converted by reduction into a corresponding amino compound, or

- 20 if a compound of general formula I is obtained wherein R₄ denotes a phenyl group substituted by an amino, alkylamino, aminoalkyl or N-alkyl-amino group, this may subsequently be converted, by reaction with a corresponding cyanate, isocyanate or carbamoyl halide, into a corresponding urea compound of general formula I, or

- 25 if a compound of general formula I is obtained wherein R₄ denotes a phenyl group substituted by an amino, alkylamino, aminoalkyl or N-alkyl-amino group, this may subsequently be converted, by reaction with a corresponding compound which transfers the amidino group or by reaction with a corresponding nitrile, into a corresponding guanidino compound of general formula I.

30

The subsequent hydrolysis is preferably carried out in an aqueous solvent, e.g. in water, methanol/water, ethanol/water, isopropanol/water, tetrahydrofuran/water or dioxan/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulphuric acid or in the presence of an alkali metal base such as lithium hydroxide, sodium hydroxide or potassium hydroxide at temperatures between 0 and 100°C, preferably at temperatures between 10 and 50°C.

The subsequent reductive alkylation is preferably carried out in a suitable solvent such as methanol, methanol/water, methanol/water/ammonia, ethanol, ether, tetrahydrofuran, dioxan or dimethylformamide, optionally with the addition of an acid such as hydrochloric acid in the presence of catalytically activated hydrogen, e.g. hydrogen in the presence of Raney nickel, platinum or palladium/charcoal, or in the presence of a metal hydride such as sodium borohydride, lithium borohydride, sodium cyanoborohydride or lithium aluminium hydride at temperatures between 0 and 100°C, preferably at temperatures between 20 and 80°C.

The subsequent acylation or sulphonylation is preferably carried out with the corresponding free acid or a corresponding reactive compound such as the anhydride, ester, imidazolide or halide thereof, preferably in a solvent such as methylene chloride, diethylether, tetrahydrofuran, toluene, dioxan, acetonitrile, dimethylsulphoxide or dimethylformamide, optionally in the presence of an inorganic or tertiary organic base at temperatures between -20 and 200°C, preferably at temperatures between 20°C and boiling temperature of the solvent used. The reaction with the free acid may optionally be carried out in the presence of an acid-activating agent or a dehydrating agent, e.g. in the presence of isobutyl chloroformate, tetraethyl orthocarbonate, trimethyl orthoacetate, 2,2-dimethoxypropane, tetramethoxysilane, thionyl chloride, trimethylchlorosilane, phosphorus trichloride, phosphorus pentoxide, N,N'-dicyclohexylcarbodiimide, N,N'-dicyclohexylcarbodiimide/N-hydroxysuccinimide, N,N'-dicyclohexylcarbodiimide/1-hydroxy-benzotriazole, 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium-tetrafluoroborate, 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium-tetrafluoroborate/1-hydroxy-benzotriazole, N,N'-carbonyldiimidazole

or triphenylphosphine/carbon tetrachloride, and optionally with the addition of a base such as pyridine, 4-dimethylamino-pyridine, N-methyl-morpholine or triethylamine, conveniently at temperatures between 0 and 150°C, preferably at temperatures between 0 and 100°C. The reaction with a corresponding reactive compound may optionally be

5 carried out in the presence of a tertiary organic base such as triethylamine, N-ethyl-diisopropylamine, N-methyl-morpholine or pyridine or by using an anhydride in the presence of the corresponding acid at temperatures between 0 and 150°C, preferably at temperatures between 50 and 100°C.

10 The subsequent esterification or amidation is conveniently carried out by reacting a corresponding reactive carboxylic acid derivative with a corresponding alcohol or amine as described hereinbefore.

The subsequent oxidation of the sulphur atom is preferably carried out in a solvent or

15 mixture of solvents, e.g. in water, water/pyridine, acetone, methylene chloride, acetic acid, acetic acid/acetic anhydride, dilute sulphuric acid or trifluoroacetic acid, usefully at temperatures of between -80 and 100°C depending on the oxidising agent used.

In order to prepare a corresponding sulphonyl compound of general formula I the oxidation

20 is expediently carried out with one equivalent of the oxidising agent used, e.g. with hydrogen peroxide in glacial acetic acid, trifluoroacetic acid or formic acid at 0 to 20°C or in acetone at 0 to 60°C, with a peracid such as performic acid in glacial acetic acid or trifluoroacetic acid at 0 to 50°C or with m-chloroperbenzoic acid in methylene chloride, chloroform or dioxan at -20 to 80°C, with sodium metaperiodate in aqueous methanol or

25 ethanol at -15 to 25°C, with bromine in glacial acetic acid or aqueous acetic acid optionally in the presence of a weak base such as sodium acetate, with N-bromosuccinimide in ethanol, with tert.butyl hypochlorite in methanol at -80 to -30°C, with iodobenzodichloride in aqueous pyridine at 0 to 50°C, with nitric acid in glacial acetic acid at 0 to 20°C, with chromic acid in glacial acetic acid or in acetone at 0 to 20°C and with sulphuryl chloride in

30 methylene chloride at -70°C, the resulting thioether-chlorine complex is expediently hydrolysed with aqueous ethanol.

In order to prepare a sulphonyl compound of general formula I the oxidation is expediently carried out starting from a corresponding sulphanyl compound with one or more equivalents of the oxidising agent used or starting from a corresponding mercapto
 5 compound, expediently with two or more equivalents of the oxidising agent used, e.g. with hydrogen peroxide in glacial acetic acid/acetic anhydride, trifluoroacetic acid or in formic acid at 20 to 100°C or in acetone at 0 to 60°C, with a peracid such as performic acid or m-chloroperbenzoic acid in glacial acetic acid, trifluoroacetic acid, methylene chloride or chloroform at temperatures between 0 and 60°C, with nitric acid in glacial acetic acid at 0
 10 to 20°C, with chromic acid, sodium periodate or potassium permanganate in acetic acid, water/sulphuric acid or in acetone at 0 to 20°C.

The subsequent reduction of a nitro group is preferably carried out by hydrogenolysis, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal or Raney nickel in a
 15 solvent such as methanol, ethanol, ethyl acetate, dimethylformamide, dimethylformamide/acetone or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid or glacial acetic acid at temperatures of between 0 and 50°C, but preferably at ambient temperature, and at a hydrogen pressure of 1 to 7 bar, but preferably 3 to 5 bar.

20 The subsequent preparation of a corresponding urea compound of general formula I is conveniently carried out with an inorganic cyanate or a corresponding isocyanate or carbamoylchloride, preferably in a solvent such as dimethylformamide and optionally in the presence of a tertiary organic base such as triethylamine at temperatures between 0 and
 25 50°C, preferably at ambient.

The subsequent preparation of a corresponding guanidino compound of general formula I is conveniently carried out by reacting with a compound which transfers the amidino group such as 3,5-dimethylpyrazole-1-carboxylic acid amidine, preferably in a solvent such as
 30 dimethylformamide and optionally in the presence of a tertiary organic base such as triethylamine at temperatures of between 0 and 50°C, preferably at ambient temperature.

In the reactions described hereinbefore, any reactive groups present such as carboxy, hydroxy, amino, alkylamino or imino groups may be protected during the reaction by
5 conventional protecting groups which are cleaved again after the reaction.

For example, a protecting group for a carboxyl group may be a trimethylsilyl, methyl, ethyl, tert.butyl, benzyl or tetrahydropyranyl group and

10 protecting groups for a hydroxy, amino, alkylamino or imino group may be an acetyl, trifluoroacetyl, benzoyl, ethoxycarbonyl, tert.butoxycarbonyl, benzyloxycarbonyl, benzyl, methoxybenzyl or 2,4-dimethoxybenzyl group and additionally, for the amino group, a phthalyl group.

15 Any protecting group used is optionally subsequently cleaved for example by hydrolysis in an aqueous solvent, e.g. in water, isopropanol/water, tetrahydrofuran/water or dioxan/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulphuric acid or in the presence of an alkali metal base such as lithium hydroxide, sodium hydroxide or potassium hydroxide, at temperatures between 0 and 100°C, preferably at temperatures
20 between 10 and 50°C.

However, a benzyl, methoxybenzyl or benzyloxycarbonyl group is cleaved, for example, hydrogenolytically, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a solvent such as methanol, ethanol, ethyl acetate,
25 dimethylformamide, dimethylformamide/acetone or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid or glacial acetic acid at temperatures between 0 and 50°C, but preferably at ambient temperature, and at a hydrogen pressure of 1 to 7 bar, but preferably 3 to 5 bar.

30 A methoxybenzyl group may also be cleaved in the presence of an oxidising agent such as cerium(IV)ammonium nitrate in a solvent such as methylene chloride, acetonitrile or

acetonitrile/water at temperatures of between 0 and 50°C, but preferably at ambient temperature.

5 A 2,4-dimethoxybenzyl group, however, is preferably cleaved in trifluoroacetic acid in the presence of anisole.

10 A tert.butyl or tert.butyloxycarbonyl group is preferably cleaved by treating with an acid such as trifluoroacetic acid or hydrochloric acid, optionally using a solvent such as methylene chloride, dioxan, ethyl acetate or ether.

15 A phthalyl group is preferably cleaved in the presence of hydrazine or a primary amine such as methylamine, ethylamine or n-butylamine in a solvent such as methanol, ethanol, isopropanol, toluene/water or dioxan at temperatures between 20 and 50°C.

20 Moreover, chiral compounds of general formula I obtained may be resolved into their enantiomers and/or diastereomers.

25 Thus, for example, the compounds of general formula I obtained which occur as racemates may be separated by methods known *per se* (cf. Allinger N. L. and Eliel E. L. in "Topics in Stereochemistry", Vol. 6, Wiley Interscience, 1971) into their optical antipodes and compounds of general formula I with at least 2 asymmetric carbon atoms may be resolved into their diastereomers on the basis of their physical-chemical differences using methods known *per se*, e.g. by chromatography and/or fractional crystallisation, and, if these compounds are obtained in racemic form, they may subsequently be resolved into the enantiomers as mentioned above.

30 The enantiomers are preferably separated by column separation on chiral phases or by recrystallisation from an optically active solvent or by reacting with an optically active substance which forms salts or derivatives such as e.g. esters or amides with the racemic compound, particularly acids and the activated derivatives or alcohols thereof, and separating the mixture of diastereomeric salts or derivatives thus obtained, e.g. on the basis

of their differences in solubility, whilst the free antipodes may be released from the pure diastereomeric salts or derivatives by the action of suitable agents. Optically active acids in common use are e.g. the D- and L-forms of tartaric acid or dibenzoyltartaric acid, di-o-tolyltartaric acid, malic acid, mandelic acid, camphorsulphonic acid, glutamic acid, N-acetylglutamic acid, aspartic acid, N-acetylaspartic acid or quinic acid. An optically active alcohol may be for example (+)- or (-)-menthol and an optically active acyl group in amides, for example, may be a (+)- or (-)-menthyloxycarbonyl group.

Furthermore, the compounds of formula I obtained may be converted into the salts thereof, particularly for pharmaceutical use into the physiologically acceptable salts with inorganic or organic acids. Acids which may be used for this purpose include for example hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, fumaric acid, succinic acid, lactic acid, citric acid, tartaric acid, maleic acid or methanesulphonic acid.

Moreover, if the new compounds of formula I thus obtained contain a carboxy group, they may subsequently, if desired, be converted into the salts thereof with inorganic or organic bases, particularly for pharmaceutical use into the physiologically acceptable salts thereof. Suitable bases for this purpose include for example sodium hydroxide, potassium hydroxide, cyclohexylamine, ethanolamine, diethanolamine and triethanolamine.

The compounds of general formulae VII to XII used as starting materials are known from the literature in some cases or may be obtained by methods known from the literature or may be obtained by the methods described hereinbefore and in the Examples. For example, the compounds of general formula VI are described in German Patent Application 198 24 922.5.

Moreover, the compounds of general formula XI may be obtained from the compounds of general formula I wherein R₄ denotes a C₁₋₄-alkyl-phenyl group substituted in the alkyl moiety by a hydroxy group, for example, by reacting with alkyl- or arylsulphonyl - chlorides.

As already mentioned, the new compounds of general formula I wherein R₁ denotes a hydrogen atom or a prodrug group have valuable pharmacological properties, particularly inhibitory effects on various kinases, especially on receptor-tyrosine kinases such as VEGFR2, PDGFR α , PDGFR β , FGFR1, FGFR3, EGFR, HER2, IGF1R and HGFR, as well as on complexes of CDK's (Cyclin Dependent Kinases) such as CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK8 and CDK9 with their specific cyclins (A, B1, B2, C, D1, D2, D3, E, F, G1, G2, H, I and K) and on viral cyclin, on the proliferation of cultivated human cells, particularly endothelial cells, e.g. in angiogenesis, but also on the proliferation of other cells, particularly tumour cells.

The biological properties of the new compounds were tested by the following standard procedure, as follows:

Human umbilical endothelial cells (HUVEC) were cultivated in IMDM (Gibco BRL), supplemented with 10 % foetal calf serum (FBS) (Sigma), 50 μ M of β -mercaptoethanol (Fluka), standard antibiotics, 15 μ g/ml of endothelial cell growth factor (ECGS, Collaborative Biomedical Products) and 100 μ g/ml of heparin (Sigma) on gelatine-coated culture dishes (0.2 % gelatine, Sigma) at 37°C, under 5 % CO₂ in a water-saturated atmosphere.

In order to investigate the inhibitory activity of the compounds according to the invention the cells were starved for 16 hours, i.e. kept in culture medium without growth factors (ECGS + heparin). The cells were detached from the culture dishes using trypsin/EDTA and washed once in serum-containing medium. Then they were seeded out in amounts of 2.5 x 10³ cells per well.

The proliferation of the cells was stimulated with 5 ng/ml of VEGF₁₆₅ (vascular endothelial growth factor; H. Weich, GBF Braunschweig) and 10 μ g/ml of heparin. As a control, 6 wells in each dish were not stimulated.

The compounds according to the invention were dissolved in 100% dimethylsulphoxide and added to the cultures in various dilutions in triplicate, the maximum dimethyl sulphoxide concentration being 0.3 %.

- 5 The cells were incubated for 76 hours at 37°C, then for a further 16 hours ³H-thymidine (0.1 µ Ci/well, Amersham) was added in order to determine the DNA synthesis. Then the radioactively labelled cells were immobilised on filter mats and the radioactivity incorporated was measured in a β-counter. In order to determine the inhibitory activity of the compounds according to the invention the mean value of the non-stimulated cells was
10 subtracted from the mean value of the factor-stimulated cells (in the presence or absence of the compounds according to the invention).

- The relative cell proliferation was calculated as a percentage of the control (HUVEC without inhibitor) and the concentration of active substance which inhibits the proliferation
15 of the cells by 50 % (IC₅₀) was determined.

The test results of the following compounds (a) to (s) of general formula I are given by way of example:

- 20 (a) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

- (b) 3-Z-[(1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone,

- 25 (c) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

- (d) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-
30 indolinone,

- (e) 3-Z-[1-(4-((2,6-dimethyl-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,
- (f) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,
- (g) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,
- (h) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,
- (i) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,
- (j) 3-Z-[1-(4-(N-acetyl-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,
- (k) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,
- (l) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,
- (m) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,
- (n) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(o) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(p) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(q) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(r) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone and

(s) 3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone.

The following Table contains the results found:

Compound	IC ₅₀ [μ m]
(a)	0.04
(b)	0.35
(c)	0.01
(d)	0.02
(e)	0.05
(f)	0.01
(g)	0.003
(h)	0.01
(i)	0.03
(j)	0.02
(k)	0.03
(l)	0.1
(m)	0.02
(n)	0.02
(o)	0.01
(p)	0.02
(q)	0.02
(r)	0.01
(s)	0.04

In view of their inhibitory effect on the proliferation of cells, particularly endothelial cells
 5 and tumour cells, the compounds of general formula I are suitable for treating diseases in
 which the proliferation of cells, particularly endothelial cells, plays a part.

Thus, for example, the proliferation of endothelial cells and the concomitant
 neovascularisation constitute a crucial stage in tumour progression (Folkman J. et al.,

Nature 339, 58-61, (1989); Hanahan D. and Folkman J., Cell 86, 353-365, (1996)).

Furthermore, the proliferation of endothelial cells is also important in haemangiomas, in metastatisation, rheumatoid arthritis, psoriasis and ocular neovascularisation (Folkman J., Nature Med. 1, 27-31, (1995)). The therapeutic usefulness of inhibitors of endothelial cell proliferation was demonstrated in the animal model for example by O'Reilly et al. and Parangi et al. (O'Reilly M.S. et al., Cell 88, 277-285, (1997); Parangi S. et al., Proc Natl Acad Sci USA 93, 2002-2007, (1996)).

The compounds of general formula I, their tautomers, their stereoisomers or the physiologically acceptable salts thereof are thus suitable, for example, for treating tumours (e.g. plate epithelial carcinoma, astrocytoma, Kaposi's sarcoma, glioblastoma, lung cancer, bladder cancer, carcinoma of the neck, melanoma, ovarian cancer, prostate cancer, breast cancer, small-cell lung cancer, glioma, colorectal carcinoma, urogenital cancer and gastrointestinal carcinoma as well as haematological cancers, such as multiple myeloma), psoriasis, arthritis (e.g. rheumatoid arthritis), haemangioma, angiofibroma, eye diseases (e.g. diabetic retinopathy), neovascular glaucoma, kidney diseases (e.g. glomerulonephritis), diabetic nephropathy, malignant nephrosclerosis, thrombic microangiopathic syndrome, transplant rejections and glomerulopathy, fibrotic diseases (e.g. cirrhosis of the liver), mesangial cell proliferative diseases, arteriosclerosis and damage to the nerve tissue and also for inhibiting the reocclusion of blood vessels after treatment with a balloon catheter, in vascular prosthetics or after the insertion of mechanical devices for keeping blood vessels open (e.g. stents), or other diseases in which cell proliferation or angiogenesis are involved.

By reason of their biological properties the compounds according to the invention may be used on their own or in conjunction with other pharmacologically active compounds, for example in tumour therapy, in monotherapy or in conjunction with other anti-tumour therapeutic agents, for example in combination with topoisomerase inhibitors (e.g. etoposide), mitosis inhibitors (e.g. vinblastin, taxol), compounds which interact with nucleic acids (e.g. cis-platin, cyclophosphamide, adriamycin), hormone antagonists (e.g.

tamoxifen), inhibitors of metabolic processes (e.g. 5-FU etc.), cytokines (e.g. interferons), kinase inhibitors, antibodies, or in conjunction with radiotherapy, etc. These combinations may be administered either simultaneously or sequentially.

- 5 For pharmaceutical use the compounds according to the invention are generally used for warm-blooded vertebrates, particularly humans, in doses of 0.01-100 mg/kg of body weight, preferably 0.1-20 mg/kg. For administration they are formulated with one or more conventional inert carriers and/or diluents, e.g. with corn starch, lactose, glucose, microcrystalline cellulose, magnesium stearate, polyvinylpyrrolidone, citric acid, tartaric
10 acid, water, water/ethanol, water/glycerol, water/sorbitol, water/polyethylene glycol, propylene glycol, stearyl alcohol, carboxymethylcellulose or fatty substances such as hard fat or suitable mixtures thereof in conventional galenic preparations such as plain or coated tablets, capsules, powders, injectable solutions, ampoules, suspensions, solutions, sprays or suppositories.

15

The Examples which follow are intended to illustrate the invention:

Abbreviations used:

20

FMOC = 9-fluorenylmethoxycarbonyl

HOBt = 1-hydroxy-1H-benzotriazole

25 TBTU = O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium-tetrafluoroborate

DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene

Preparation of the starting compounds:

Solid phase Example I

- 5 2.0 g of Rink resin (MBHA resin, made by Messrs Novabiochem) are left to swell in 30 ml of dimethylformamide. Then 40 ml of 30% piperidine in dimethylformamide are added and the mixture is shaken for 7 minutes to cleave the Fmoc protecting group. The resin is then washed repeatedly with dimethylformamide. Then 0.4 g of 2-indolinone-6-carboxylic acid (prepared analogously to Langenbeck et al., Justus Liebigs Ann. Chem. 499, 201-208 (1932)), 297 mg HOBt, 706 mg TBTU and 0.9 ml of N-ethyl-diisopropylamine in 30 ml of dimethylformamide are added and the mixture is shaken for 1 hour. Then the solution is suction filtered and the resin is washed five times with 30 ml of dimethylformamide and three times with 30 ml of methylene chloride. To dry it, nitrogen is blown through the resin.
- 10
- 15 Yield: 1.9 g of charged resin

Solid phase example II

- 1.9 g of the resin obtained in Example I are stirred with 6 ml of acetic anhydride and 6 ml of triethyl orthobenzoate for 3 hours at 110°C. Then the mixture is left to cool and the resin is washed with dimethylformamide and subsequently with methylene chloride.
- 20
- Yield: 1.9 g of moist resin

The following charged resins are prepared analogously to Example II:

- 25
- (1) resin charged with 3-Z-(1-ethoxy-methylene)-6-carbamoyl-2-indolinone
Prepared by reacting the resin obtained according to Example I with triethyl orthoformate
- (2) resin charged with 3-Z-(1-methoxy-1-methyl-methylene)-6-carbamoyl-2-indolinone
- 30 Prepared by reacting the resin obtained according to Example I with trimethyl orthoacetate

(3) resin charged with 3-Z-(1-methoxy-1-ethyl-methylene)-6-carbamoyl-2-indolinone
Prepared by reacting the resin obtained according to Example I with trimethyl
orthopropionate

- 5 (4) resin charged with 3-Z-(1-methoxy-1-propyl-methylene)-6-carbamoyl-2-indolinone
Prepared by reacting the product of Example I and trimethyl orthobutyrate

Example III

- 10 N-(4-nitrophenyl)-N-methyl-methanesulphonamide
-

3.0 g of N-methyl-4-nitroaniline are dissolved in 20 ml of pyridine and 2.4 g of
methanesulphonic acid chloride added dropwise at room temperature. The mixture is
stirred for 12 hours at room temperature. After this time the mixture is poured onto water,
15 the precipitate formed is filtered off and dried at 50°C *in vacuo*.
Yield: 4.0 g (87 % of theory),
R_f value: 0.5 (silica gel, ethyl acetate/toluene = 7:3)
Melting point: 107-108 °C

- 20 Example IV

N-(2-dimethylamino-ethyl)-N-methylsulphonyl-4-nitroaniline

- 38.9 g of N-methylsulphonyl-4-nitroaniline are dissolved in 2.0 l of acetone, 51.9 g of 1-
25 chloro-2-dimethylamino-ethane, 77.4 g of potassium carbonate and 5.0 g of sodium iodide
are added and the mixture is stirred for a total of 4 days at 50°C, while after 12 hours a
further 25.9 g of 1-chloro-2-dimethylamino-ethane, 49.8 g of potassium carbonate and 5.0
g of sodium iodide in 500 ml of acetone are added and after 36 hours another 26.0 g of 1-
chloro-2-dimethylamino-ethane, 50.0 g of potassium carbonate and 5.0 g of sodium iodide
30 in 100 ml of acetone are added. After this time the mixture is filtered and the filtrate
evaporated down. The residue is stirred with ether, suction filtered and dried at 40°C.

Yield: 25.3 g (49 % of theory),

R_f value: 0.5 (silica gel, methylene chloride/methanol/ammonia = 9:1:0.1)

C₁₁H₁₇N₃O₄S

ESI mass spectrum: m/z = 288 [M+H⁺]

5

The following compounds are prepared analogously to Example IV:

(1) 4-[N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino]-nitrobenzene

10 (2) N-carboxymethyl-N-methylsulphonyl-4-nitroaniline

(3) N-cyanomethyl-N-methylsulphonyl-p-phenylenediamine

(4) 4-[N-(2-(N-benzyl-N-methyl-amino)-ethyl)-N-methylsulphonyl-amino]-nitrobenzene

15

(5) 4-[N-(3-phthalimido-2-yl-propyl)-N-methylsulphonyl-amino]-nitrobenzene

(6) 4-[N-(3-(N-benzyl-N-methyl-amino)-propyl)-N-methylsulphonyl-amino]-nitrobenzene

20 Example V

N-(dimethylaminocarbonyl-methyl)-N-methylsulphonyl-4-nitroaniline

7.0 g of N-carboxymethyl-N-methylsulphonyl-4-nitroaniline, 2.5 g of dimethylamine
25 hydrochloride, 8.1 g of TBTU and 3.9 g of HOBT are dissolved in 125 ml of
dimethylformamide and at 0°C 17.6 ml of N-ethyl-diisopropylamine are added. The
mixture is stirred for 4 hours at room temperature, diluted with 1 l of water and the
precipitate formed is suction filtered. After washing with water, ethanol and ether the
residue is dried at 70°C *in vacuo*.

30 Yield: 5.3 g (69 % of theory),

R_f value: 0.40 (silica gel, methylene chloride/methanol = 9:1)



ESI mass spectrum: $m/z = 300$ [M-H]

The following compounds are prepared analogously to Example V:

5

(1) 4-[(N-dimethylaminocarbonylmethyl)-amino]-nitrobenzene

prepared from 4-(N-carboxymethyl-amino)-nitrobenzene and dimethylamine hydrochloride

10

(2) 4-(N-methylaminocarbonylmethyl-N-methylsulphonyl-amino)-nitrobenzene

Prepared from N-carboxymethyl-N-methylsulphonyl-4-nitroaniline and methylamine hydrochloride

(3) 4-[(N-(methylcarbamoyl-methyl)-N-methyl-amino)-methyl]-nitrobenzene

15

Prepared from 4-[(N-carboxymethyl-N-methyl-amino)-methyl]-nitrobenzene and methylamine hydrochloride

(4) 4-[(N-(dimethylcarbamoyl-methyl)-N-methyl-amino)-methyl]-nitrobenzene

20

Prepared from 4-[(N-carboxymethyl-N-methyl-amino)-methyl]-nitrobenzene and dimethylamine hydrochloride

Example VI

4-[N-(2-dimethylamino-ethyl)-N-acetyl-amino]-nitrobenzene

25

3.6 g of 4-(2-dimethylamino-ethylamino)-nitrobenzene (according to Gabbay et al., J. Am. Chem. Soc. 91, 5136 (1969)) are dissolved in 50 ml of methylene chloride and 5.0 ml of triethylamine are added. 1.3 ml of acetyl chloride are slowly added dropwise to this mixture at room temperature and the mixture is stirred for 2 hours at room temperature.

30

After this time another 5.0 ml of triethylamine and 1.3 ml of acetylchloride are added and the mixture is refluxed for another 2 hours. The solvent is removed, the residue is taken up

in ethyl acetate and the organic phase is extracted twice with water. After drying over MgSO_4 the solvent is removed and the residue dried *in vacuo*.

Yield: 2.0 g (45 % of theory),

R_f value: 0.55 (silica gel, methylene chloride/methanol/ammonia = 9:1:0.1)

5 $\text{C}_{12}\text{H}_{17}\text{N}_3\text{O}_3$

ESI mass spectrum: $m/z = 252$ $[\text{M}+\text{H}^+]$

The following compounds are prepared analogously to Example VI:

10 (1) 4-[N-(3-dimethylamino-propyl)-N-acetyl-amino]-nitrobenzene

Prepared from 4-(3-dimethylamino-propylamino)-nitrobenzene (according to Gabbay et al., J. Am. Chem. Soc. 91, 5136 (1969) and acetyl chloride

(2) 4-[N-(2-dimethylamino-ethyl)-N-propionyl-amino]-nitrobenzene

15 Prepared from 4-(2-dimethylamino-ethylamino)-nitrobenzene and propionyl chloride

(3) 4-[N-acetyl-N-(dimethylaminocarbonylmethyl)-amino]-nitrobenzene

Prepared from 4-[N-(dimethylaminocarbonylmethyl)-amino]-nitrobenzene and acetyl chloride

20

(4) 4-[N-(2-dimethylamino-ethyl)-N-butyryl-amino]-nitrobenzene

Prepared from 4-(2-dimethylamino-ethylamino)-nitrobenzene and butyryl chloride

(5) 4-[N-(2-dimethylamino-ethyl)-N-isobutyryl-amino]-nitrobenzene

25 Prepared from 4-(2-dimethylamino-ethylamino)-nitrobenzene and isobutyryl chloride

(6) 4-[N-(2-dimethylamino-ethyl)-N-benzoyl-amino]-nitrobenzene

Prepared from 4-(2-dimethylamino-ethylamino)-nitrobenzene and benzoyl chloride

30 (7) 4-[N-(2-dimethylamino-ethyl)-N-acetyl-amino]-1,3-dinitrobenzene

Prepared from 4-(2-dimethylamino-ethyl-amino)-1,3-dinitrobenzene and acetyl chloride

(8) 4-[N-(2-dimethylamino-ethyl)-N-(furan-2-carbonyl)-amino]-nitrobenzene

Prepared from 4-(2-dimethylamino-ethylamino)-nitrobenzene and furan-2-carbonyl chloride

5

(9) 4-[N-(2-dimethylamino-ethyl)-N-(2-methoxy-benzoyl)-amino]-nitrobenzene

Prepared from 4-(2-dimethylamino-ethylamino)-nitrobenzene and 2-methoxy-benzoyl chloride

10 (10) 4-[N-(2-dimethylamino-ethyl)-N-(pyridine-3-carbonyl)-amino]-nitrobenzene

Prepared from 4-(2-dimethylamino-ethylamino)-nitrobenzene and nicotinic acid chloride

(11) 4-[N-(2-dimethylamino-ethyl)-N-(phenyl-acetyl)-amino]-nitrobenzene

Prepared from 4-(2-dimethylamino-ethylamino)-nitrobenzene and phenylacetyl-chloride

15

(12) 4-[N-(2-dimethylamino-ethyl)-N-acetyl-amino]-3-bromo-nitrobenzene

Prepared from 4-[N-(2-dimethylamino-ethyl)-amino]-3-bromo-nitrobenzene and acetyl chloride

20 (13) N-acryloyl-N-methyl-4-nitro-aniline

Prepared from 4-methylamino-nitrobenzene and acrylic acid chloride

(14) N-acryloyl-N-isopropyl-4-nitro-aniline

Prepared from 4-isopropylamino-nitrobenzene and acrylic acid chloride

25

(15) N-acryloyl-N-benzyl-4-nitro-aniline

Prepared from 4-benzylamino-nitrobenzene and acrylic acid chloride

(16) N-bromoacetyl-N-methyl-4-nitro-aniline

30 Prepared from 4-methylamino-nitrobenzene and bromoacetyl chloride

(17) N-bromoacetyl-N-isopropyl-4-nitro-aniline

Prepared from 4-isopropylamino-nitrobenzene and bromoacetyl chloride

(18) N-bromoacetyl-N-benzyl-4-nitro-aniline

5 Prepared from 4-benzylamino-nitrobenzene and bromoacetyl chloride

Example VII

N-(dimethylaminomethylcarbonyl)-N-methyl-4-nitro-aniline

10

1.8 g of dimethylamine hydrochloride and 5.5 g of potassium carbonate are placed in 80 ml of acetone and 4.2 g of N-bromoacetyl-N-methyl-4-nitroaniline are added in three batches at room temperature. The mixture is stirred for 12 hours at room temperature. After this time the mixture is filtered and the filtrate is evaporated down. The residue is dissolved in
15 ethyl acetate, washed twice with water, dried over sodium sulphate and finally concentrated by rotary evaporation.

Yield: 2.8 g (79 % of theory),

R_f value: 0.5 (silica gel, ethyl acetate/methanol = 7:3)

Melting point: 121-122°C

20

The following compounds are prepared analogously to Example VII:

(1) N-(piperidin-1-yl-methylcarbonyl)-N-methyl-4-nitroaniline

25 (2) N-(morpholin-4-yl-methylcarbonyl)-N-methyl-4-nitroaniline

(3) N-[(4-benzyl-piperazin-1-yl)-methylcarbonyl]-N-methyl-4-nitroaniline

(4) N-(pyrrolidin-1-yl-methylcarbonyl)-N-methyl-4-nitroaniline

30

- (5) N-[(N-aminocarbonylmethyl-N-methyl-amino)-methylcarbonyl]-N-methyl-4-nitroaniline
- (6) N-[(N-benzyl-N-methyl-amino)-methylcarbonyl]-N-methyl-4-nitroaniline
- 5 (7) N-[di-(2-methoxyethyl)-amino-methylcarbonyl]-N-methyl-4-nitroaniline
- (8) N-(dimethylaminomethylcarbonyl)-N-isopropyl-4-nitro-aniline
- 10 (9) N-(piperidin-1-yl-methylcarbonyl)-N-isopropyl-4-nitro-aniline
- (10) N-[(4-tert.butoxycarbonyl-piperazin-1-yl)-methylcarbonyl]-N-isopropyl-4-nitro-aniline
- 15 (11) N-[(N-benzyl-N-methyl-amino)-methylcarbonyl]-N-benzyl-4-nitro-aniline
- (12) N-(dimethylaminomethylcarbonyl)-N-benzyl-4-nitro-aniline
- (13) N-(piperidin-1-yl-methylcarbonyl)-N-benzyl-4-nitro-aniline
- 20 (14) N-[di-(2-hydroxyethyl)-amino-methylcarbonyl]-N-methyl-4-nitroaniline
- (15) N-[(N-(2-methoxyethyl)-N-methyl-amino)-methylcarbonyl]-N-methyl-4-nitroaniline
- 25 (16) N-[(N-(2-dimethylamino-ethyl)-N-methyl-amino)-methylcarbonyl]-N-methyl-4-nitroaniline
- (17) N-[(4-methyl-piperazin-1-yl)-methylcarbonyl]-N-methyl-4-nitroaniline
- 30 (18) N-[(imidazol-1-yl)-methylcarbonyl]-N-methyl-4-nitroaniline

(19) N-[(phthalimido-2-yl)-methylcarbonyl]-N-methyl-4-nitroaniline

Example VIII

5 N-[(2-dimethylamino-ethyl)-carbonyl]-N-benzyl-4-nitro-aniline

0.5 g of dimethylamine hydrochloride, 1.1 ml of triethylamine and 1.2 g of N-acryloyl-N-benzyl-4-nitro-aniline are dissolved in 50 ml of methanol and stirred for 24 hours at room temperature. After this time the mixture is evaporated down. The residue is purified over
 10 an aluminium oxide column (activity 2-3) with methylene chloride/ethanol 50:1 as eluant.
 Yield: 1.4 g (98 % of theory),
 R_f value: 0.8 (aluminium oxide, methylene chloride/ethanol = 20:1)
 Melting point: 73°C

15 The following compounds are prepared analogously to Example VIII:

(1) N-[(2-dimethylamino-ethyl)-carbonyl]-N-isopropyl-4-nitro-aniline

Prepared from N-acryloyl-N-isopropyl-4-nitro-aniline and dimethylamine hydrochloride

20 (2) N-[(2-dimethylamino-ethyl)-carbonyl]-N-methyl-4-nitro-aniline

Prepared from N-acryloyl-N-methyl-4-nitro-aniline and dimethylamine hydrochloride

(3) N-[(2-(4-tert.butoxycarbonyl-piperazin-1-yl)-ethyl)-carbonyl]-N-methyl-4-nitro-aniline

Prepared from N-acryloyl-N-methyl-4-nitro-aniline and N-tert.butoxycarbonyl-piperazine

25

(4) N-[(2-(piperidin-1-yl)-ethyl)-carbonyl]-N-methyl-4-nitro-aniline

Prepared from N-acryloyl-N-methyl-4-nitro-aniline and piperidine

(5) N-[(2-(N-benzyl-N-methyl-amino)-ethyl)-carbonyl]-N-methyl-4-nitro-aniline

30 Prepared from N-acryloyl-N-methyl-4-nitro-aniline and N-benzyl-N-methyl-amine

Example IX

4-(4-methyl-piperazine-1-yl)-nitrobenzene

- 5 31.5 g of 4-chloro-1-nitrobenzene and 44.4 ml of 1-methylpiperazine are combined and stirred for 18 hours at 90°C. Then the solution is poured onto ice water and the precipitate formed is suction filtered, washed with water and recrystallised from ethanol/water 1:1. The residue is dried *in vacuo* at 75°C.
Yield: 44.0 g (99 % of theory),
10 R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)
Melting point: 108-112°C

The following compounds are prepared analogously to Example IX:

- 15 (1) N-(2-dimethylaminoethyl)-N-methyl-4-nitroaniline
Prepared from 1-fluoro-4-nitrobenzene and 1-dimethylamino-2-methylamino-ethane

 (2) N-(3-dimethylaminopropyl)-N-methyl-4-nitroaniline
Prepared from 1-fluoro-4-nitrobenzene and 1-dimethylamino-3-methylamino-propane
20 (3) 4-(N-carboxymethyl-amino)-nitrobenzene
Prepared from 1-fluoro-4-nitrobenzene and glycine

 (4) N-cyclohexyl-p-phenylenediamine
25 Prepared from 1-fluoro-4-nitrobenzene and cyclohexylamine

 (5) 6-[N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino]-3-phthalimido-2-yl-nitrobenzene
Prepared from 2-nitro-4-phthalimido-2-yl-fluorobenzene,
30 N-(2-dimethylamino-ethyl)-methanesulphonamide and sodium hydride as base

(6) 6-[N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino]-1,3-dinitrobenzene

Prepared from 2,4-dinitro-chlorobenzene, N-(2-dimethylamino-ethyl)-methanesulphonamide and sodium hydride as base

5 (7) 4-[N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino]-3-chloro-nitrobenzene

Prepared from 2-fluoro-5-nitro-chlorobenzene, N-(2-dimethylamino-ethyl)-methanesulphonamide and sodium hydride as base

(8) 4-(2-dimethylamino-ethyl-amino)-1,3-dinitrobenzene

10 Prepared from 1-chloro-2,4-dinitro-benzene and N,N-dimethyl-ethylenediamine

(9) 4-[N-(2-dimethylamino-ethyl)-N-(ethylsulphonyl)-amino]-nitrobenzene

Prepared from 1-fluoro-4-nitro-benzene, N-(2-dimethylamino-ethyl)-ethanesulphonamide and sodium hydride as base

15

(10) 4-[N-(2-dimethylamino-ethyl)-N-(propylsulphonyl)-amino]-nitrobenzene

Prepared from 1-fluoro-4-nitro-benzene, N-(2-dimethylamino-ethyl)-propanesulphonamide and sodium hydride as base

20 (11) 4-[N-(2-dimethylamino-ethyl)-N-(butylsulphonyl)-amino]-nitrobenzene

Prepared from 1-fluoro-4-nitro-benzene, N-(2-dimethylamino-ethyl)-butanesulphonamide and sodium hydride as base

(12) 4-[N-(2-dimethylamino-ethyl)-N-(benzylsulphonyl)-amino]-nitrobenzene

25 Prepared from 1-fluoro-4-nitro-benzene, N-(2-dimethylamino-ethyl)-C-phenylmethanesulphonamide and sodium hydride as base

(13) 4-[N-(2-dimethylamino-ethyl)-N-(phenylsulphonyl)-amino]-nitrobenzene

30 Prepared from 1-fluoro-4-nitro-benzene, N-(2-dimethylamino-ethyl)-benzenesulphonamide and sodium hydride as base

(14) 4-[N-(2-dimethylamino-ethyl)-N-(isopropylsulphonyl)-amino]-nitrobenzene

Prepared from 1-fluoro-4-nitro-benzene, N-(2-dimethylamino-ethyl)-isopropylsulphonamide and sodium hydride as base

5 (15) 4-[N-(2-dimethylamino-ethyl)-amino]-3-bromo-nitrobenzene

Prepared from 2-bromo-1-fluoro-4-nitro-benzene and N,N-dimethyl-ethylenediamine

(16) 4-isopropylamino-nitrobenzene

Prepared from 1-fluoro-4-nitrobenzene and isopropylamine

10

(17) 4-benzylamino-nitrobenzene

Prepared from 1-fluoro-4-nitrobenzene and benzylamine

Example X

15

4-(imidazol-4-yl)-nitrobenzene

20

9.5 g of 2-phenylimidazole are carefully dissolved in 50 ml of concentrated sulphuric acid and 5.8 g of ammonium nitrate are added to this solution at 0°C. After a further 60 minutes stirring at 0°C the mixture is poured onto ice water, made basic with ammonia water and the precipitate formed is suction filtered and recrystallised from ethanol.

Yield: 8.0 g (64 % of theory),

R_f value: 0.6 (silica gel, ethyl acetate/ethanol = 10:1)

C₉H₇N₃O₂

25

Mass spectrum: m/z = 189 [M⁺]

The following compounds are prepared analogously to Example X:

(1) 4-(imidazol-2-yl)-nitrobenzene

30

Prepared from 4-(imidazol-2-yl)-benzene

(2) 4-(5-methyl-imidazol-4-yl)-nitrobenzene

Prepared from 4-methyl-5-phenyl-imidazole (J. Heterocycl. Chem. 1983, 20, 1277-1281)

Example XI

5

4-(2-(imidazol-4-yl)-ethylene)-nitrobenzene

1.5 g of 4-nitrobenzaldehyde and 7.45 g of (N-trityl-imidazol-4-yl-methyl)-
triphenylphosphonium chloride are dissolved in 75 ml of tetrahydrofuran and to this
10 solution 3.0 ml of DBU are added dropwise at room temperature. After a further 120
minutes stirring at room temperature the mixture is poured onto water and the precipitate
formed is suction filtered. The product is taken up in 25 ml of 1N hydrochloric acid and
refluxed for 4 hours. After this time it is neutralised with ammoniacal water, extracted with
ethyl acetate and the organic phase is washed with water, dried over sodium sulphate and
15 evaporated down. The residue is purified over a silica gel column with methylene
chloride/methanol 10:1 as eluant.
Yield: 1.0 g of (47 % of theory),
R_f value: 0.6 (silica gel, ethyl acetate/ethanol = 10:1)
Melting point: 185-188°C

20

Example XII

4-(piperidin-1-yl-methyl)-nitrobenzene

25 40.0 g of 4-nitrobenzyl bromide are dissolved in 500 ml of methylene chloride, 51.5 ml of
triethylamine are added and 18.3 ml of piperidine are carefully added dropwise. After the
end of the exothermic reaction the mixture is refluxed for another 30 minutes. After
cooling it is washed with water and the organic phase is dried over sodium sulphate.
Finally, the organic phase is evaporated down.
30 Yield: 36.3 g of (89 % of theory),
R_f value: 0.6 (silica gel, methylene chloride/methanol = 9:1)



Mass spectrum: $m/z = 221$ [M^+]

The following compounds are prepared analogously to Example XII:

5

(1) 4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-nitrobenzene

(2) 3-(N,N-dimethyl-aminomethyl)-nitrobenzene

10

(3) 4-(N,N-dimethyl-aminomethyl)-nitrobenzene

(4) 4-(2-dimethylamino-ethyl)-nitrobenzene

(5) 4-(2-diethylamino-ethyl)-nitrobenzene

15

(6) 4-(diethylamino-methyl)-nitrobenzene

(7) 4-(N-benzyl-N-methyl-aminomethyl)-nitrobenzene

20

(8) 4-(N-ethyl-N-methyl-aminomethyl)-nitrobenzene

(9) 4-[N-(n-hexyl)-N-methyl-aminomethyl]-nitrobenzene

(10) 4-(thiomorpholin-4-yl-methyl)-nitrobenzene

25

(11) 4-[(4-methyl-piperazine-1-yl)-methyl]-nitrobenzene

(12) 4-(imidazol-1-yl-methyl)-nitrobenzene

30

(13) 4-[2-(4-hydroxy-piperidin-1-yl)-ethyl-amino]-nitrobenzene

- (14) 4-[(3-hydroxy-pyrrolidin-1-yl)-methyl]-nitrobenzene
- (15) 4-(1,2,4-triazol-1-yl-methyl)-nitrobenzene
- 5 (16) 4-(1,2,3-triazol-2-yl-methyl)-nitrobenzene
- (17) 4-(1,2,3-triazol-1-yl-methyl)-nitrobenzene
- (18) 4-[(N-ethoxycarbonylmethyl-N-methyl-amino)-methyl]-nitrobenzene
- 10 (19) 4-[(N-aminocarbonylmethyl-N-methyl-amino)-methyl]-nitrobenzene
- (20) 4-(azetidin-1-yl-methyl)-nitrobenzene
- 15 (21) 4-[(di-(2-methoxy-ethyl)-amino)-methyl]-nitrobenzene
- (22) 4-[N-(N-tert.butoxycarbonyl-3-amino-propyl)-N-methyl-aminomethyl]-nitrobenzene
- (23) 4-[(N-propyl-N-methyl-amino)-methyl]-nitrobenzene
- 20 (24) 4-[(N-(2-dimethylamino-ethyl)-N-methyl-amino)-methyl]-nitrobenzene
- (25) 4-[(N-(3-dimethylamino-propyl)-N-methyl-amino)-methyl]-nitrobenzene
- 25 (26) 4-[(N-(2-methoxy-ethyl)-N-methyl-amino)-methyl]-nitrobenzene
- (27) 4-[(N-(2-hydroxy-ethyl)-N-methyl-amino)-methyl]-nitrobenzene
- (28) 4-[(N-(dioxolan-2-yl-methyl)-N-methyl-amino)-methyl]-nitrobenzene
- 30 (29) 4-(3-oxo-piperazine-1-yl-methyl)-nitrobenzene

Example XIII4-[(N-carboxymethyl-N-methyl-amino)-methyl]-nitrobenzene

5

7.33 g of 4-[(N-ethoxycarbonylmethyl-N-methyl-amino)-methyl]-nitrobenzene are dissolved in 140 ml of ethanol, 34.0 ml of 1N sodium hydroxide solution are added and the mixture is stirred for half an hour at room temperature. After this time the mixture is neutralised with 34 ml of 1N hydrochloric acid, the solvent removed, the residue taken up in methylene chloride and extracted with water. The aqueous phase is evaporated down and the residue is recrystallised from methylene chloride.

10

Yield: 5.43 g (84 % of theory),

R_f value: 0.4 (silica gel, methylene chloride/methanol = 2:1)

C₁₀H₁₂N₂O₄

15

Mass spectrum: m/z = 223 [M⁺]

Example XIV4-(N-ethyl-aminomethyl)-nitrobenzene

20

6.0 g of 4-nitrobenzyl bromide are dissolved in 25 ml of ethanol, combined with 25 ml of 10% ethanolic ethylamine solution and refluxed for 2 hours. Then the solution is concentrated by rotary evaporation, the residue is taken up with methylene chloride and washed with dilute sodium hydroxide solution. Finally the organic phase is evaporated down.

25

Yield: 2.3 g (46 % of theory),

R_f value: 0.2 (silica gel, methylene chloride/methanol = 9:1)

C₉H₁₂N₂O₂

ESI mass spectrum: m/z = 179 [M-H⁻]

30

The following compounds are prepared analogously to Example XIV:

- (1) 4-[N-(4-chlorobenzyl)-aminomethyl]-nitrobenzene
- (2) 4-(N-cyclohexyl-aminomethyl)-nitrobenzene
- 5 (3) 4-(N-isopropyl-aminomethyl)-nitrobenzene
- (4) 4-(N-propyl-aminomethyl)-nitrobenzene
- 10 (5) 4-(N-methyl-aminomethyl)-nitrobenzene
- (6) 4-(N-butyl-aminomethyl)-nitrobenzene
- (7) 4-(N-methoxycarbonylmethyl-aminomethyl)-nitrobenzene
- 15 (8) 4-(N-benzyl-aminomethyl)-nitrobenzene
- (9) 4-(aminomethyl)-nitrobenzene
- 20 (10) 4-(pyrrolidin-1-yl-methyl)-nitrobenzene
- (11) 4-(morpholin-4-yl-methyl)-nitrobenzene
- (12) 4-(hexamethyleneiminomethyl)-nitrobenzene
- 25 (13) 4-(4-hydroxy-piperidin-1-yl-methyl)-nitrobenzene
- (14) 4-(4-methoxy-piperidin-1-yl-methyl)-nitrobenzene
- 30 (15) 4-(4-methyl-piperidin-1-yl-methyl)-nitrobenzene

- (16) 4-(4-ethyl-piperidin-1-yl-methyl)-nitrobenzene
- (17) 4-(4-isopropyl-piperidin-1-yl-methyl)-nitrobenzene
- 5 (18) 4-(4-phenyl-piperidin-1-yl-methyl)-nitrobenzene
- (19) 4-(4-benzyl-piperidin-1-yl-methyl)-nitrobenzene
- (20) 4-(4-ethoxycarbonyl-piperidin-1-yl-methyl)-nitrobenzene
- 10 (21) 4-(N,N-dipropyl-aminomethyl)-nitrobenzene
- (22) 4-(4-tert.butoxycarbonyl-piperazin-1-yl-methyl)-nitrobenzene
- 15 (23) 4-(2-morpholin-4-yl-ethyl)-nitrobenzene
- (24) 4-(2-pyrrolidin-1-yl-ethyl)-nitrobenzene
- (25) 4-(2-piperidin-1-yl-ethyl)-nitrobenzene
- 20 (26) 4-(N-ethyl-N-benzyl-aminomethyl)-nitrobenzene
- (27) 4-(N-propyl-N-benzyl-aminomethyl)-nitrobenzene
- 25 (28) 4-[N-methyl-N-(4-chlorobenzyl)-aminomethyl]-nitrobenzene
- (29) 4-[N-methyl-N-(4-bromobenzyl)-aminomethyl]-nitrobenzene
- (30) 4-[N-methyl-N-(4-fluorobenzyl)-aminomethyl]-nitrobenzene
- 30 (31) 4-[N-methyl-N-(4-methylbenzyl)-aminomethyl]-nitrobenzene

- (32) 4-[N-methyl-N-(3-chlorobenzyl)-aminomethyl]-nitrobenzene
- (33) 4-[N-methyl-N-(3,4-dimethoxybenzyl)-aminomethyl]-nitrobenzene
- 5 (34) 4-[N-methyl-N-(4-methoxybenzyl)-aminomethyl]-nitrobenzene
- (35) 4-(N-2,2,2-trifluoroethyl-N-benzyl-aminomethyl)-nitrobenzene
- 10 (36) 4-[N-2,2,2-trifluoroethyl-N-(4-chlorobenzyl)-aminomethyl]-nitrobenzene
- (37) 4-(thiomorpholin-4-yl-methyl)-nitrobenzene
- (38) 4-(azetidin-1-yl-methyl)-nitrobenzene
- 15 (39) 4-(3,4-dihydropyrrolidin-1-yl-methyl)-nitrobenzene
- (40) 4-(3,4-dihydropiperidin-1-yl-methyl)-nitrobenzene
- 20 (41) 4-(2-methoxycarbonyl-pyrrolidin-1-yl-methyl)-nitrobenzene
- (42) 4-(3,5-dimethyl-piperidin-1-yl-methyl)-nitrobenzene
- (43) 4-(4-phenyl-piperazin-1-yl-methyl)-nitrobenzene
- 25 (44) 4-(4-phenyl-4-hydroxy-piperidin-1-yl-methyl)-nitrobenzene
- (45) 4-[N-(3,4,5-trimethoxybenzyl)-N-methyl-aminomethyl]-nitrobenzene
- 30 (46) 4-[N-(3,4-dimethoxybenzyl)-N-ethyl-aminomethyl]-nitrobenzene

- (47) 4-[N-(2,6-dichlorobenzyl)-N-methyl]-aminomethyl]-nitrobenzene
- (48) 4-[N-(4-trifluoromethylbenzyl)-N-methyl]-aminomethyl]-nitrobenzene
- 5 (49) 4-(N-benzyl-N-isopropyl-aminomethyl)-nitrobenzene
- (50) 4-(N-benzyl-N-tert.butyl-aminomethyl)-nitrobenzene
- (51) 4-(N,N-diisopropyl-aminomethyl)-nitrobenzene
- 10 (52) 4-(N,N-diisobutyl-aminomethyl)-nitrobenzene
- (53) 4-(2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-nitrobenzene
- 15 (54) 4-(2,3-dihydro-isoindol-2-yl-methyl)-nitrobenzene
- (55) 4-(6,7-dimethoxy-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-nitrobenzene
- (56) 4-(1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-nitrobenzene
- 20 (57) 4-[N-(2-hydroxyethyl)-N-benzyl-aminomethyl]-nitrobenzene
- (58) 4-[N-(1-ethyl-pentyl)-N-(pyridin-2-yl-methyl)-aminomethyl]-nitrobenzene
- 25 (59) 4-(piperin-1-yl-methyl)-1,3-dinitrobenzene
- (60) 4-(N-phenethyl-N-methyl-aminomethyl)-nitrobenzene
- (61) 4-[N-(3,4-dihydroxy-phenethyl)-N-methyl-aminomethyl]-nitrobenzene
- 30 (62) 4-[N-(3,4,5-trimethoxy-phenethyl)-N-methyl-aminomethyl]-nitrobenzene

- (63) 4-[N-(3,4-dimethoxy-phenethyl)-N-methyl-aminomethyl]-nitrobenzene
- (64) 4-[N-(3,4-dimethoxy-benzyl)-N-methyl-aminomethyl]-nitrobenzene
- 5 (65) 4-[N-(4-chloro-benzyl)-N-methyl-aminomethyl]-nitrobenzene
- (66) 4-[N-(4-bromo-benzyl)-N-methyl-aminomethyl]-nitrobenzene
- 10 (67) 4-[N-(4-fluoro-benzyl)-N-methyl-aminomethyl]-nitrobenzene
- (68) 4-[N-(4-methyl-benzyl)-N-methyl-aminomethyl]-nitrobenzene
- (69) 4-[N-(4-nitro-phenethyl)-N-methyl-aminomethyl]-nitrobenzene
- 15 (70) 4-(N-phenethyl-N-benzyl-aminomethyl)-nitrobenzene
- (71) 4-(N-phenethyl-N-cyclohexyl-aminomethyl)-nitrobenzene
- 20 (72) 4-[N-(2-(pyridin-2-yl)-ethyl)-N-methyl-aminomethyl]-nitrobenzene
- (73) 4-[N-(2-(pyridin-4-yl)-ethyl)-N-methyl-aminomethyl]-nitrobenzene
- (74) 4-[N-(pyridin-4-yl-methyl)-N-methyl-aminomethyl]-nitrobenzene
- 25 (75) 4-(N,N-dibenzyl-aminomethyl)-nitrobenzene
- (76) 4-[N-(4-nitro-phenethyl)-N-propyl-aminomethyl]-nitrobenzene
- 30 (77) 4-(N-benzyl-N-(3-cyano-propyl)-aminomethyl)-nitrobenzene

- (78) 4-(N-benzyl-N-allyl-aminomethyl)-nitrobenzene
- (79) 4-[N-benzyl-N-(2,2,2-trifluoroethyl)-aminomethyl]-nitrobenzene
- 5 (80) 4-[N-(2-benzo(1,3)dioxol-5-yl-methyl)-N-methyl-aminomethyl]-nitrobenzene
- (81) 4-(7-chloro-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-nitrobenzene
- (82) 4-(7,8-dichloro-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-nitrobenzene
- 10 (83) 4-(7-methoxy-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-nitrobenzene
- (84) 4-(7-methyl-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-nitrobenzene
- 15 (85) 4-(7,8-dimethoxy-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-nitrobenzene
- (86) 4-(6,7-dichloro-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-nitrobenzene
- (87) 4-(6,7-dimethyl-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-nitrobenzene
- 20 (88) 4-(6-chloro-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-nitrobenzene
- (89) 4-(7-chloro-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-nitrobenzene
- 25 (90) 4-(6-methoxy-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-nitrobenzene
- (91) 4-(7-methoxy-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-nitrobenzene
- (92) 4-[(2,3,4,5-tetrahydro-azepino(4,5-b)pyrazin-3-yl)-methyl]-nitrobenzene
- 30 (93) 4-[(7-amino-2,3,4,5-tetrahydro-azepino(4,5-b)pyrazin-3-yl)-methyl]-nitrobenzene

(94) 4-[(2-amino-5,6,7,8-tetrahydro-azepino(4,5-d)thiazol-6-yl)-methyl]-nitrobenzene

(95) 4-[(5,6,7,8-tetrahydro-azepino(4,5-d)thiazol-6-yl)-methyl]-nitrobenzene

5

Example XV

4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-nitrobenzene

- 10 6.0 g of 4-(thiomorpholin-4-yl-methyl)-nitrobenzene are dissolved in 100 ml of methylene chloride and 10.3 g of meta-chloroperbenzoic acid are slowly added. After a further 3 hours stirring at room temperature the precipitate obtained is filtered off.

Yield: 6.2 g (91 % of theory)

R_f value: 0.5 (silica gel, methylene chloride/methanol = 1:1)

- 15 C₁₁H₁₄N₂O₄S

Mass spectrum: m/z = 270 [M⁺]

The following compound is prepared analogously to Example XV:

- 20 (1) 4-(1-oxo-thiomorpholin-4-yl-methyl)-nitrobenzene

Example XVI

4-[N-(3-amino-propyl)-N-methylsulphonyl-amino]-nitrobenzene

25

9.5 g of 4-[N-(3-phthalimido-2-yl-propyl)-N-methylsulphonyl-amino]-nitrobenzene are dissolved in 200 ml of ethanol, 11.5 ml of hydrazine hydrate are added and the mixture is stirred for 1.5 hours at 50 °C. After cooling the residue is largely evaporated down, water is added and the solution is extracted with methylene chloride. The organic phase is dried, 30 evaporated down and purified over a silica gel column with methylene chloride/methanol/ammonia 9:1:0.1.

Yield: 2.5 g (39 % of theory)

R_f value: 0.2 (silica gel, methylene chloride/methanol = 9:1)

C₁₀H₁₅N₃O₄S

ESI mass spectrum: m/z = 272 [M-H]⁻

5

The following compound is prepared analogously to Example XVI:

(1) 6-[N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino]-3-amino-nitrobenzene

Prepared from 6-[N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino]-3-phthalimido-2-yl-nitrobenzene

10

Example XVII

4-(1-methyl-imidazol-2-yl)-nitrobenzene

15

7.5 g of 4-(imidazol-2-yl)-nitrobenzene are dissolved in 50 ml of dimethylsulphoxide and at 0°C 5.0 g of potassium tert.butoxide are added. After one hour of stirring at room temperature 2.6 ml of methyl iodide are added dropwise and the mixture is stirred for one hour at room temperature. After this time the residue is poured onto ice water and the precipitate formed is suction filtered, washed with water and dried.

20

Yield: 6.1 g (76 % of theory)

R_f value: 0.6 (silica gel, methylene chloride/methanol = 10:1)

Melting point: 186-187°C

25 The following compounds are prepared analogously to Example XVII:

(1) 4-(1-ethyl-imidazol-2-yl)-nitrobenzene

Prepared from 4-(imidazol-2-yl)-nitrobenzene and ethyl iodide

30 (2) 4-(1-benzyl-imidazol-2-yl)-nitrobenzene

Prepared from 4-(imidazol-2-yl)-nitrobenzene and benzyl bromide

Example XVIII

4-[(N-(2-(2-methoxy-ethoxy)-ethyl)-N-methyl-amino)-methyl]-nitrobenzene

- 5 5.0 g of 4-methylaminomethyl-nitrobenzene are dissolved in 30 ml of dimethylformamide and 4.6 g of 2-(2-methoxy-ethoxy)-ethyl chloride are added. After six hours' stirring at 100°C the solvent is removed and the residue is taken up in ethyl acetate. The organic phase is washed with water and dried over sodium sulphate. After the elimination of the solvent the residue is purified over an aluminium oxide column (activity 2-3) with
- 10 toluene/ethyl acetate 5:1 as eluant.

Yield: 2.3 g (29 % of theory)

R_f value: 0.5 (aluminium oxide, toluene/ethyl acetate = 5:1)

C₁₃H₂₀N₂O₄

ESI mass spectrum: m/z = 267 [M-H⁻]

15

Example XIX

4-(N-ethyl-N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

- 20 2.2 g of 4-(ethylaminomethyl)-nitrobenzene are dissolved in 50 ml of ethyl acetate and stirred with 2.6 g of di-tert-butyl. dicarbonate (tert.butoxycarbonyl-anhydride) for 30 minutes at room temperature. Then the solution is washed with water and evaporated down.

Yield: 3.4 g of theory

- 25 R_f value: 0.3 (silica gel, methylene chloride/methanol = 50:1)

Melting point: 85 °C

The following compounds are prepared analogously to Example XIX:

- 30 (1) 4-[N-(4-chlorophenyl-methyl)-N-tert.butoxycarbonyl-aminomethyl]-nitrobenzene

(2) 4-(N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

(3) 4-(N-cyclohexyl-N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

5 (4) 4-(N-isopropyl-N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

(5) 4-(N-methyl-N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

(6) 4-(N-propyl-N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

10

(7) 4-(N-butyl-N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

(8) 4-(N-methoxycarbonylmethyl-N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

15 (9) 4-(N-benzyl-N-tert.butoxycarbonyl-aminomethyl)-nitrobenzene

(10) 4-[N-(3-trifluoroacetyl-amino-propyl)-N-methylsulphonyl-amino]-nitrobenzene

Prepared from 4-[N-(3-amino-propyl)-N-methylsulphonyl-amino]-nitrobenzene and trifluoroacetic acid anhydride

20

(11) 4-[(4-tert.butoxycarbonyl-piperazin-1-yl)-methyl]-nitrobenzene

Example XX

25 4-(piperidin-1-yl-methyl)-aniline

37.0 g of 4-(piperidin-1-yl-methyl)-nitrobenzene are dissolved in 300 ml of methanol, 8.0 g of Raney nickel are added and the mixture is hydrogenated for 85 minutes with 3 bars of hydrogen at room temperature. The catalyst is filtered off and the filtrate is evaporated

30 down.

Yield: 24.0 g (75 % of theory),

R_f value: 0.4 (silica gel, methylene chloride/methanol = 9:1)

C₁₂H₁₈N₂

ESI mass spectrum: m/z = 191 [M+H⁺]

5 The following compounds are prepared analogously to Example VIII:

(1) 4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-aniline

(2) N-(2-dimethylamino-ethyl)-N-methylsulphonyl-p-phenylenediamine

10

(3) 3-(dimethylaminomethyl)-aniline

(4) 4-(dimethylaminomethyl)-aniline

15

(5) 4-(2-dimethylamino-ethyl)-aniline

(6) 4-[N-(2-dimethylamino-ethyl)-N-acetyl-amino]-aniline

(7) 4-[N-(3-dimethylamino-propyl)-N-acetyl-amino]-aniline

20

(8) 4-[N-(2-dimethylamino-ethyl)-N-benzoyl-amino]-aniline

(9) 4-[N-(2-dimethylamino-ethyl)-N-propionyl-amino]-aniline

25

(10) 4-[N-(2-dimethylamino-ethyl)-N-butyryl-amino]-aniline

(11) 4-[N-(2-dimethylamino-ethyl)-N-isobutyryl-amino]-aniline

(12) 4-(N-tert.butoxycarbonyl-aminomethyl)-aniline

30

(13) 4-(N-ethyl-N-tert.butoxycarbonyl-aminomethyl)-aniline

- (14) 4-[N-(4-chlorophenyl-methyl)-N-tert.butoxycarbonyl-aminomethyl]-aniline
- (15) 4-(N-cyclohexyl-N-tert.butoxycarbonyl-aminomethyl)-aniline
- 5 (16) 4-(N-isopropyl-N-tert.butoxycarbonyl-aminomethyl)-aniline
- (17) 4-(N-propyl-N-tert.butoxycarbonyl-aminomethyl)-aniline
- 10 (18) 4-(N-methyl-N-tert.butoxycarbonyl-aminomethyl)-aniline
- (19) 4-(N-butyl-N-tert.butoxycarbonyl-aminomethyl)-aniline
- (20) 4-(N-methoxycarbonyl-methyl-N-tert.butoxycarbonyl-aminomethyl)-aniline
- 15 (21) 4-(N-benzyl-N-tert.butoxycarbonyl-aminomethyl)-aniline
- (22) 4-(pyrrolidin-1-yl-methyl)-aniline
- 20 (23) 4-(morpholin-4-yl-methyl)-aniline
- (24) 4-(hexamethyleneiminomethyl)-aniline
- (25) 4-(4-hydroxy-piperidin-1-yl-methyl)-aniline
- 25 (26) 4-(4-methoxy-piperidin-1-yl-methyl)-aniline
- (27) 4-(4-methyl-piperidin-1-yl-methyl)-aniline
- 30 (28) 4-(4-ethyl-piperidin-1-yl-methyl)-aniline

(29) 4-(4-isopropyl-piperidin-1-yl-methyl)-aniline

(30) 4-(4-phenyl-piperidin-1-yl-methyl)-aniline

5 (31) 4-(4-benzyl-piperidin-1-yl-methyl)-aniline

(32) 4-(4-ethoxycarbonyl-piperidin-1-yl-methyl)-aniline

(33) 4-(N,N-dipropyl-aminomethyl)-aniline

10

(34) 4-(4-tert.butoxycarbonyl-piperazin-1-yl-methyl)-aniline

(35) 4-(2-morpholin-4-yl-ethyl)-aniline

15 (36) 4-(2-pyrrolidin-1-yl-ethyl)-aniline

(37) 4-(2-piperidin-1-yl-ethyl)-aniline

(38) 4-(N-propyl-N-benzyl-aminomethyl)-aniline

20

(39) 4-[N-(n-hexyl)-N-methyl-aminomethyl]-aniline

(40) 4-[N-methyl-N-(4-chlorobenzyl)-aminomethyl]-aniline

25 (41) 4-[N-methyl-N-(4-bromobenzyl)-aminomethyl]-aniline

(42) 4-[N-methyl-N-(4-methylbenzyl)-aminomethyl]-aniline

(43) 4-[N-methyl-N-(4-fluorobenzyl)-aminomethyl]-aniline

30

(44) 4-[N-methyl-N-(3-chlorobenzyl)-aminomethyl]-aniline

- (45) 4-[N-methyl-N-(3,4-dimethoxybenzyl)-aminomethyl]-aniline
- (46) 4-[N-methyl-N-(4-methoxybenzyl)-aminomethyl]-aniline
- 5 (47) 4-(N-2,2,2-trifluoroethyl-N-benzyl-aminomethyl)-aniline
- (48) 4-[N-2,2,2-trifluoroethyl-N-(4-chlorobenzyl)-aminomethyl]-aniline
- 10 (49) 4-(thiomorpholin-4-yl-methyl)-aniline
- (50) 4-(1-oxo-thiomorpholin-4-yl-methyl)-aniline
- (51) 4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-aniline
- 15 (52) 4-(azetidion-1-yl-methyl)-aniline
- (53) 4-(3,4-dihydropyrrolidin-1-yl-methyl)-aniline
- 20 (54) 4-(3,4-dihydropiperidin-1-yl-methyl)-aniline
- (55) 4-(2-methoxycarbonyl-pyrrolidin-1-yl-methyl)-aniline
- (56) 4-(3,5-dimethyl-piperidin-1-yl-methyl)-aniline
- 25 (57) 4-(4-phenyl-piperazin-1-yl-methyl)-aniline
- (58) 4-(4-phenyl-4-hydroxy-piperidin-1-yl-methyl)-aniline
- 30 (59) 4-[N-(3,4,5-trimethoxy-benzyl)-N-methyl-aminomethyl]-aniline

- (60) 4-[N-(3,4-dimethoxy-benzyl)-N-ethyl-aminomethyl]-aniline
- (61) 4-(N-benzyl-N-ethyl-aminomethyl)-aniline
- 5 (62) 4-[N-(2,6-dichlorobenzyl)-N-methyl-aminomethyl]-aniline
- (63) 4-[N-(4-trifluoromethylbenzyl)-N-methyl-aminomethyl]-aniline
- (64) 4-(N-benzyl-N-isopropyl-aminomethyl)-aniline
- 10 (65) 4-(N-benzyl-N-tert.butyl-aminomethyl)-aniline
- (66) 4-(diethylamino-methyl)-aniline
- 15 (67) 4-(2-diethylamino-ethyl)-aniline
- (68) 4-(N,N-diisopropyl-aminomethyl)-aniline
- (69) 4-(N,N-diisobutyl-aminomethyl)-aniline
- 20 (70) 4-(2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-aniline
- (71) 4-(2,3-dihydro-isoindol-2-yl-methyl)-aniline
- 25 (72) 4-(6,7-dimethoxy-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-aniline
- (73) 4-(1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-aniline
- (74) 4-[N-(2-hydroxy-ethyl)-N-benzyl-aminomethyl]-aniline
- 30 (75) 4-[N-(1-ethyl-pentyl)-N-(pyridin-2-yl-methyl)-aminomethyl]-aniline

- (76) 4-(piperidin-1-yl-methyl)-3-nitro-aniline
- (77) 4-(piperidin-1-yl-methyl)-3-amino-aniline
- 5 (78) 4-(N-benzyl-N-methyl-aminomethyl)-aniline
- (79) 4-(N-ethyl-N-methyl-aminomethyl)-aniline
- 10 (80) 4-(N-phenethyl-N-methyl-aminomethyl)-aniline
- (81) 4-[N-(3,4-dihydroxy-phenethyl)-N-methyl-aminomethyl]-aniline
- (82) 4-[N-(3,4,5-trimethoxy-phenethyl)-N-methyl-aminomethyl]-aniline
- 15 (83) 4-[N-(3,4-dimethoxy-phenethyl)-N-methyl-aminomethyl]-aniline
- (84) 4-[N-(3,4-dimethoxy-benzyl)-N-methyl-aminomethyl]-aniline
- 20 (85) 4-[N-(4-chloro-benzyl)-N-methyl-aminomethyl]-aniline
- (86) 4-[N-(4-bromo-benzyl)-N-methyl-aminomethyl]-aniline
- (87) 4-[N-(4-fluoro-benzyl)-N-methyl-aminomethyl]-aniline
- 25 (88) 4-[N-(4-methyl-benzyl)-N-methyl-aminomethyl]-aniline
- (89) 4-[N-(4-nitro-phenethyl)-N-methyl-aminomethyl]-aniline
- 30 (90) 4-(N-phenethyl-N-benzyl-aminomethyl)-aniline

(91) 4-(N-phenethyl-N-cyclohexyl-aminomethyl)-aniline

(92) 4-[N-(2-(pyridin-2-yl)-ethyl)-N-methyl-aminomethyl]-aniline

5 (93) 4-[N-(2-(pyridin-4-yl)-ethyl)-N-methyl-aminomethyl]-aniline

(94) 4-[N-(pyridin-4-yl-methyl)-N-methyl-aminomethyl]-aniline

(95) 4-(N,N-dibenzylaminomethyl)-aniline

10

(96) 4-[N-(4-nitro-benzyl)-N-propyl-aminomethyl]-aniline

(97) 4-[N-benzyl-N-(3-cyano-propyl)-aminomethyl]-aniline

15 (98) 4-(N-benzyl-N-allyl-aminomethyl)-aniline

(99) 4-[N-benzyl-N-(2,2,2-trifluoroethyl)-aminomethyl]-aniline

(100) 4-[(benzo(1,3)dioxol-5-yl-methyl)-methyl-aminomethyl]-aniline

20

(101) 4-(7-chloro-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-aniline

(102) 4-(7,8-dichloro-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-aniline

25 (103) 4-(7-methoxy-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-aniline

(104) 4-(7-methyl-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-aniline

(105) 4-(7,8-dimethoxy-2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-aniline

30

(106) 4-(6,7-dichloro-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-aniline

- (107) 4-(6,7-dimethyl-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-aniline
- (108) 4-(6-chloro-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-aniline
- 5 (109) 4-(7-chloro-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-aniline
- (110) 4-(6-methoxy-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-aniline
- 10 (111) 4-(7-methoxy-1,2,3,4-tetrahydro-isoquinolin-2-yl-methyl)-aniline
- (112) 4-(2,3,4,5-tetrahydro-azepino(4,5-b)pyrazin-3-yl-methyl)-aniline
- (113) 4-(7-amino-2,3,4,5-tetrahydro-azepino(4,5-b)pyrazin-3-yl-methyl)-aniline
- 15 (114) 4-(2-amino-5,6,7,8-tetrahydro-azepino(4,5-d)thiazol-6-yl-methyl)-aniline
- (115) 4-(5,6,7,8-tetrahydro-azepino(4,5-d)thiazol-6-yl-methyl)-aniline
- 20 (116) 4-(4-methyl-piperazin-1-yl)-aniline
- (117) 4-[N-(2-dimethylamino-ethyl)-N-methyl-amino]-aniline
- (118) 4-[N-(3-dimethylamino-propyl)-N-methyl-amino]-aniline
- 25 (119) N-(3-dimethylamino-propyl)-N-methylsulphonyl-p-phenylenediamine
- (120) 4-[(N-dimethylaminocarbonylmethyl-N-methylsulphonyl)-amino]-aniline
- 30 (121) N-(4-aminophenyl)-N-methyl-methanesulphonamide

(122) 4-(imidazol-4-yl)-aniline

(123) 4-(tetrazol-5-yl)-aniline

5 (124) 4-[N-(2-dimethylamino-ethyl)-N-propionyl-amino]-aniline

(125) N-(dimethylaminomethylcarbonyl)-N-methyl-p-phenylenediamine

(126) N-[(2-dimethylamino-ethyl)-carbonyl]-N-methyl-p-phenylenediamine

10

(127) 4-(N-acetyl-N-dimethylaminocarbonylmethyl)-amino)-aniline

(128) N-methylaminocarbonylmethyl-N-methylsulphonyl-p-phenylenediamine

15 (129) N-aminocarbonylmethyl-N-methylsulphonyl-p-phenylenediamine

(130) 4-(imidazolidin-2,4-dion-5-ylidene-methyl)-aniline

(131) 4-(imidazolidin-2,4-dion-5-yl-methyl)-aniline

20

(132) 4-(2-oxo-pyrrolidin-1-yl-methyl)-aniline

(133) N-cyanomethyl-N-methylsulphonyl-p-phenylenediamine

25 (134) 4-[2-(imidazol-4-yl)-ethyl]-aniline

(135) 4-[(4-methyl-piperazin-1-yl)-methyl]-aniline

(136) 4-[N-(2-(N-benzyl-N-methyl-amino)-ethyl)-N-methylsulphonyl-amino]-aniline

30

(137) 4-[N-(3-(N-benzyl-N-methyl-amino)-propyl)-N-methylsulphonyl-amino]-aniline

- (138) N-cyclohexyl-p-phenylenediamine
- (139) 4-(pyridin-4-yl-methyl)-aniline
- 5 (140) 4-(imidazol-1-yl-methyl)-aniline
- (141) 4-benzyl-aniline
- 10 (142) N-(3-trifluoroacetyl-amino-propyl)-N-methylsulphonyl-p-phenylenediamine
- (143) tert.butyl 4-amino-phenylacetate
- (144) 4-(imidazol-2-yl)-aniline
- 15 (145) 4-(1-methyl-imidazol-2-yl)-aniline
- (146) 4-(1-ethyl-imidazol-2-yl)-aniline
- 20 (147) 4-(1-benzyl-imidazol-2-yl)-aniline
- (148) 4-[N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino]-3-amino-aniline
- (149) 4-[N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino]-3-chloro-aniline
- 25 (150) 4-[N-(2-dimethylamino-ethyl)-N-acetyl-amino]-3-amino-aniline
- (151) 4-[N-(2-dimethylamino-ethyl)-N-acetyl-amino]-3-bromo-aniline
- 30 (152) 4-[2-(4-hydroxy-piperidin-1-yl)-ethyl-amino]-aniline

- (153) N-(2-dimethylamino-ethyl)-N-ethylsulphonyl-p-phenylenediamine
- (154) N-(2-dimethylamino-ethyl)-N-propylsulphonyl-p-phenylenediamine
- 5 (155) N-(2-dimethylamino-ethyl)-N-isopropylsulphonyl-p-phenylenediamine
- (156) N-(2-dimethylamino-ethyl)-N-butylsulphonyl-p-phenylenediamine
- (157) N-(2-dimethylamino-ethyl)-N-benzylsulphonyl-p-phenylenediamine
- 10 (158) N-(2-dimethylamino-ethyl)-N-phenylsulphonyl-p-phenylenediamine
- (159) 4-((3-hydroxy-pyrrolidin-1-yl)-methyl)-aniline
- 15 (160) 4-[N-(2-dimethylamino-ethyl)-N-(furan-2-carbonyl)-amino]-aniline
- (161) 4-[N-(2-dimethylamino-ethyl)-N-(2-methoxy-benzoyl)-amino]-aniline
- (162) 4-[N-(2-dimethylamino-ethyl)-N-(pyridine-3-carbonyl)-amino]-aniline
- 20 (163) 4-[N-(2-dimethylamino-ethyl)-N-(phenyl-acetyl)-amino]-aniline
- (164) N-(piperidin-1-yl-methylcarbonyl)-N-methyl-p-phenylenediamine
- 25 (165) N-(morpholin-4-yl-methylcarbonyl)-N-methyl-p-phenylenediamine
- (166) N-[(4-benzyl-piperazin-1-yl)-methylcarbonyl]-N-methyl-p-phenylenediamine
- (167) N-(pyrrolidin-1-yl-methylcarbonyl)-N-methyl-p-phenylenediamine
- 30 (168) 4-(5-methyl-imidazol-4-yl)-aniline

- (169) N-[(2-dimethylamino-ethyl)-carbonyl]-N-isopropyl-p-phenylenediamine
- (170) N-[(2-dimethylamino-ethyl)-carbonyl]-N-benzyl-p-phenylenediamine
- 5 (171) N-(N-aminocarbonylmethyl-N-methyl-amino)-methylcarbonyl)-N-methyl-p-phenylenediamine
- (172) N-[(N-benzyl-N-methyl-amino)-methylcarbonyl]-N-methyl-p-phenylenediamine
- 10 (173) N-[di-(2-methoxyethyl)-amino-methylcarbonyl]-N-methyl-p-phenylenediamine
- (174) N-[(2-(4-tert.butoxycarbonyl-piperazin-1-yl)-ethyl)-carbonyl]-N-methyl-p-phenylenediamine
- 15 (175) N-[(2-(piperidin-1-yl)-ethyl)-carbonyl]-N-methyl-p-phenylenediamine
- (176) N-[(2-(N-benzyl-N-methyl-amino)-ethyl)-carbonyl]-N-methyl-p-phenylenediamine
- 20 (177) N-(dimethylaminomethylcarbonyl)-N-isopropyl-p-phenylenediamine
- (178) N-(piperidin-1-yl-methylcarbonyl)-N-isopropyl-p-phenylenediamine
- (179) N-[(4-tert.butoxycarbonyl-piperazin-1-yl)-methylcarbonyl]-N-isopropyl-p-phenylenediamine
- 25 (180) N-[(N-benzyl-N-methyl-amino)-methylcarbonyl]-N-benzyl-p-phenylenediamine
- (181) N-(dimethylaminomethylcarbonyl)-N-benzyl-p-phenylenediamine
- 30 (182) N-(piperidin-1-yl-methylcarbonyl)-N-benzyl-p-phenylenediamine

- (183) 4-(1,2,4-triazol-1-yl-methyl)-aniline
- (184) 4-(1,2,3-triazol-2-yl-methyl)-aniline
- 5 (185) 4-(1,2,3-triazol-1-yl-methyl)-aniline
- (186) 4-[(N-ethoxycarbonylmethyl-N-methyl-amino)-methyl]-aniline
- 10 (187) 4-[(N-aminocarbonylmethyl-N-methyl-amino)-methyl]-aniline
- (188) 4-(azetidin-1-yl-methyl)-aniline
- (189) 4-[(di-(2-methoxy-ethyl)-amino)-methyl]-aniline
- 15 (190) 4-[(N-(2-(2-methoxy-ethoxy)-ethyl)-N-methyl-amino)-methyl]-aniline
- (191) 4-[N-(N-tert.butoxycarbonyl-3-amino-propyl)-N-methyl-aminomethyl]-aniline
- 20 (192) 4-[(N-(methylcarbamoyl-methyl)-N-methyl-amino)-methyl]-aniline
- (193) 4-[(N-(dimethylcarbamoyl-methyl)-N-methyl-amino)-methyl]-aniline
- (194) 4-[(N-propyl-N-methyl-amino)-methyl]-aniline
- 25 (195) 4-[(N-(2-dimethylamino-ethyl)-N-methyl-amino)-methyl]-aniline
- (196) 4-[(N-(3-dimethylamino-propyl)-N-methyl-amino)-methyl]-aniline
- 30 (197) 4-[(N-(2-methoxy-ethyl)-N-methyl-amino)-methyl]-aniline

(198) 4-[(N-(2-hydroxy-ethyl)-N-methyl-amino)-methyl]-aniline

(199) 4-[(N-(dioxolan-2-yl-methyl)-N-methyl-amino)-methyl]-aniline

5 (200) 4-(3-oxo-piperazin-1-yl-methyl)-aniline

(201) N-[di-(2-hydroxyethyl)-amino-methylcarbonyl]-N-methyl-p-phenylenediamine

(202) N-[(N-(2-methoxyethyl)-N-methyl-amino)-methylcarbonyl]-N-methyl-p-
10 phenylenediamine

(203) N-[(N-(2-dimethylamino-ethyl)-N-methyl-amino)-methylcarbonyl]-N-methyl-p-
phenylenediamine

15 (204) N-[(4-methyl-piperazin-1-yl)-methylcarbonyl]-N-methyl-p-phenylenediamine

(205) N-[(imidazol-1-yl)-methylcarbonyl]-N-methyl-p-phenylenediamine

(206) N-[(phthalimido-2-yl)-methylcarbonyl]-N-methyl-p-phenylenediamine
20

Example XXI

4-(4-hydroxymethyl-piperidin-1-yl-methyl-amino)-aniline

25 1.1 g of 4-(4-ethoxycarbonyl-piperidin-1-yl-methyl-amino)-aniline are suspended in 15 ml
of tetrahydrofuran. 175 mg of lithium borohydride are added at room temperature, stirred
for 24 h, another 175 mg of lithium borohydride are added and after a further 7.5 hours 15
ml of water are added and the mixture is stirred for 10 minutes. It is extracted three times
with 15 ml of ethyl acetate. The combined organic phases are washed with water and
30 saturated saline solution, dried over sodium sulphate and concentrated by rotary

evaporation. The residue is purified over a silica gel column with methylene chloride/methanol/ammonia 4:1:0.01 as eluant.

Yield: 200 mg (27 % of theory)

R_f value: 0.4 (silica gel, methylene chloride/methanol/ammonia 4:1:0.01)

5 Melting point: 157°C

Example XXII

methyl 4-methoxycarbonylmethyl-3-nitro-benzoate

10

54.3 g of methyl 3-nitro-benzoate and 29.0 g of methyl chloroacetate are dissolved in 100 ml of dimethylformamide and this solution is added dropwise at -10°C to a solution of 78.5 g of potassium-tert. butoxide in 500 ml of dimethylformamide. The mixture is stirred for another 10 minutes at room temperature and after this time the solution is poured onto 350 ml of concentrated hydrochloric acid in 2 l of ice water. The solution is stirred for 0.5 hours, the precipitate obtained is suction filtered and washed with water. The product is recrystallised from 150 ml of methanol and dried at 40°C *in vacuo*.

15

Yield: 48.3 g of (51 % of theory), contains about 20 % of methyl 6-methoxycarbonylmethyl-3-nitro-benzoate

20

R_f value: 0.7 (silica gel, petroleum ether/ethyl acetate = 1:1)

Melting point: 65-73 °C

The following compound is prepared analogously to Example XXII:

25

(1) ethyl 4-methoxycarbonylmethyl-3-nitro-benzoate

Prepared from ethyl 4-thoxycarbonylmethyl-3-nitro-benzoate

Example XXIIImethyl 2-indolinone-6-carboxylate

5 48.3 g of methyl 4-methoxycarbonylmethyl-3-nitro-benzoate are dissolved in 800 ml of concentrated acetic acid, 5.0 g of palladium on carbon (10%) are added and the solution is hydrogenated for 2.5 hours at room temperature and 50 psi. The catalyst is filtered off and the filtrate is evaporated down. The residue is taken up in 150 ml of tert.-butylmethyl ether, filtered again and dried *in vacuo* at 100°C.

10 Yield: 28.6 g (98 % of theory),

R_f value: 0.4 (silica gel, methylene chloride/methanol = 10:1)

Melting point: 208-211 °C

The following compound is prepared analogously to Example XXIII:

15

(1) ethyl 2-indolinone-6-carboxylate

Prepared from ethyl 4-methoxycarbonylmethyl-3-nitro-benzoate

Example XXIV

20

1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone

15.0 g of ethyl 2-indolinone-6-carboxylate, 49.6 ml of triethyl orthobenzoate and 150 ml of acetic anhydride are stirred for 4 hours at 110°C. After this time the solvent is removed, the
25 residue is recrystallised from petroleum ether and dried *in vacuo* at 50°C.

Yield: 16.9 g (61 % of theory),

R_f value: 0.5 (silica gel, petroleum ether/methylene chloride/ethyl acetate = 5:4:1)

Melting point: 98-100°C

C₂₂H₂₁NO₅

30

The following compounds are prepared analogously to Example XXIV:

(1) 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone

Prepared from methyl 2-indolinone-6-carboxylate, triethyl orthobenzoate and acetic anhydride

5

(2) 1-acetyl-3-(1-ethoxy-1-ethyl-methylene)-6-ethoxycarbonyl-2-indolinone

Prepared from ethyl 2-indolinone-6-carboxylate, triethyl orthopropionate and acetic anhydride

10 Preparation of the final compounds:

Example 1

15 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

300 mg of resin obtained according to Example II are suspended in 3 ml of dimethylformamide and shaken with 0.2 g of 4-(piperidin-1-yl-methyl)-aniline for 22 hours at 70°C. Then it is filtered off and the resin is washed several times with methylene chloride, methanol and dimethylformamide. Then 1 ml of methanolic ammonia is added for 2 hours in order to eliminate the acetyl group. Then after further washing 4 ml of 10% trifluoroacetic acid in methylene chloride are added during another 60 minutes, the resin is separated off and the solution is evaporated down.

Yield: 69 mg

25 R_f value: 0.1 (silica gel, methylene chloride/methanol = 9:1)

$C_{28}H_{28}N_4O_2$


Mass spectrum: $m/z = 452 (m^+)$

The following compounds are prepared analogously to Example 1:

30

(1) 3-Z-(1-Anilino-1-phenyl-methylene)-6-carbamoyl-2-indolinone


Prepared from the resin obtained according to Example II and aniline



Mass spectrum: $m/z = 355$ (m^+)

- 5 (2) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-dimethylaminomethyl-aniline




- 10 Mass spectrum: $m/z = 412$ (m^+)

- (3) 3-Z-[1-(4-(2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-(2-diethylamino-ethyl)-aniline

- 15




Mass spectrum: $m/z = 454$ (m^+)

- (4) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

- 20

Prepared from the resin obtained according to Example II and 4-(morpholin-4-yl-methyl)-aniline

R_f value: 0.50 (silica gel, methylene chloride/methanol = 4:1)




- 25 Mass spectrum: $m/z = 454$ (m^+)

- (5) 3-Z-[1-(4-(1-oxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-(1-oxo-thiomorpholin-4-yl-methyl)-aniline

- 30

R_f value: 0.30 (silica gel, methylene chloride/methanol = 9:1)




Mass spectrum: $m/z = 486$ (m^+)

- 5 (6) 3-Z-[1-(4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-aniline

R_f value: 0.30 (silica gel, methylene chloride/methanol = 9:1)




- 10 Mass spectrum: $m/z = 502$ (m^+)

- (7) 3-Z-[1-(4-(benzylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

- 15 Prepared from the resin obtained according to Example II and 4-[N-(phenyl-methyl)-N-tert.butoxycarbonyl-aminomethyl]-aniline

R_f value: 0.40 (silica gel, methylene chloride/methanol = 4:1)



Mass spectrum: $m/z = 474$ (m^+)

- 20 (8) 3-Z-[1-(4-(amino-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-(N-tert.butoxycarbonyl-aminomethyl)-aniline

R_f value: 0.10 (silica gel, methylene chloride/methanol = 4:1)

- 25 $\text{C}_{23}\text{H}_{20}\text{N}_4\text{O}_2$

Mass spectrum: $m/z = 384$ (m^+)

- (9) 3-Z-[1-(4-(2,6-dimethylpiperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

- 30 Prepared from the resin obtained according to Example II and 4-(2,6-dimethylpiperidin-1-yl-methyl)-aniline

R_f value: 0.45 (silica gel, methylene chloride/methanol = 4:1)

C₃₀H₃₂N₄O₂

Mass spectrum: m/z = 480 (m⁺)

- 5 (10) 3-Z-[1-(4-(pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-(pyrrolidin-1-yl-methyl)-aniline

R_f value: 0.15 (silica gel, methylene chloride/methanol = 4:1)

- 10 C₂₇H₂₆N₄O₂

Mass spectrum: m/z = 438 (m⁺)

(11) 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

- 15 Prepared from the resin obtained according to Example II and 3-dimethylaminomethyl-aniline

R_f value: 0.23 (silica gel, methylene chloride/methanol = 4:1)

C₂₅H₂₄N₄O₂

Mass spectrum: m/z = 412 (m⁺)

- 20

(12) 3-Z-[1-(3-(N-methyl-N-ethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 3-(N-methyl-N-ethyl-aminomethyl)-aniline

- 25 R_f value: 0.23 (silica gel, methylene chloride/methanol = 4:1)

C₂₆H₂₆N₄O₂

Mass spectrum: m/z = 426 (m⁺)

- 30 (13) 3-Z-[1-(3-(methylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-(N-tert.butoxycarbonyl-N-methyl-aminomethyl)-aniline

R_f value: 0.06 (silica gel, methylene chloride/methanol = 4:1)

C₂₄H₂₂N₄O₂

5 Mass spectrum: m/z = 399 (m+H⁺)

(14) 3-Z-[1-(3-hydroxymethyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and 3-amino-benzyl alcohol

R_f value: 0.7 (silica gel, methylene chloride/methanol = 4:1)

10 C₂₃H₁₉N₃O₃

Mass spectrum: m/z = 385 (m⁺)

(15) 3-Z-[1-(4-(methoxycarbonylmethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

15 Prepared from the resin obtained according to Example II and
4-(N-methoxycarbonylmethyl-N-tert.butoxycarbonyl-aminomethyl)-aniline

R_f value: 0.40 (silica gel, methylene chloride/methanol = 9:1)

C₂₆H₂₄N₄O₄

Mass spectrum: m/z = 457 (m+H⁺)

20

(16) 3-Z-[1-(4-(N-methylsulphonyl-N-(dimethylaminocarbonyl-methyl)-amino)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and 4-(N-methylsulphonyl-N-(dimethylaminocarbonylmethyl)-amino)-aniline

25 R_f value: 0.40 (silica gel, methylene chloride/methanol = 9:1)

C₂₇H₂₇N₅O₅S

Mass spectrum: m/z = 533 (m⁺)

30

(17) 3-Z-[1-(4-(N-acetyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and 4-(N-acetyl-aminomethyl)-aniline

R_f value: 0.70 (silica gel, methylene chloride/methanol = 4:1)

C₂₅H₂₂N₄O₃

5 Mass spectrum: m/z = 426 (m⁺)

(18) 3-Z-[1-(3,4-dimethoxy-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and 3,4-dimethoxy-aniline

R_f value: 0.40 (silica gel, methylene chloride/methanol = 9:1)

10 C₂₄H₂₁N₃O₄

Mass spectrum: m/z = 415 (m⁺)

(19) 3-Z-[1-(4-(morpholin-4-yl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

15 Prepared from the resin obtained according to Example II and 4-morpholin-4-yl-aniline

R_f value: 0.20 (silica gel, methylene chloride/methanol = 9:1)

C₂₆H₂₄N₄O₃

Mass spectrum: m/z = 440 (m⁺)

20 (20) 3-Z-[1-(4-acetylamino-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and 4-acetylamino-aniline

R_f value: 0.25 (silica gel, methylene chloride/methanol = 9:1)

C₂₄H₂₀N₄O₃

Mass spectrum: m/z = 412 (m⁺)

25

(21) 3-Z-[1-(4-amino-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and 4-amino-aniline

R_f value: 0.40 (silica gel, methylene chloride/methanol = 9:1)

C₂₂H₁₈N₄O₂

30 Mass spectrum: m/z = 370 (m⁺)

(22) 3-Z-[1-(4-N-methyl-N-acetyl-amino-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and 4-(N-methyl-N-acetyl-amino)-aniline

5 $C_{25}H_{22}N_4O_3$

Mass spectrum: $m/z = 426 (m^+)$

(23) 3-Z-[1-(4-ethoxycarbonyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and ethyl 4-amino-benzoate

10 $C_{25}H_{21}N_3O_4$

Mass spectrum: $m/z = 427 (m^+)$

(24) 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and 4-amino-benzoic acid

15 R_f value: 0.11 (silica gel, methylene chloride/methanol = 9:1)

$C_{23}H_{17}N_3O_4$

Mass spectrum: $m/z = 398 (m-H^+)$

(25) 3-Z-[1-(4-benzylcarbamoyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

20 Prepared from the resin obtained according to Example II and 4-amino-benzoic acid-benzylamide

R_f value: 0.21 (silica gel, methylene chloride/methanol = 9:1)

$C_{30}H_{24}N_4O_3$

Mass spectrum: $m/z = 488 (m^+)$

25

(26) 3-Z-[1-(cyclohexyl-amino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II and cyclohexylamine

R_f value: 0.60 (silica gel, methylene chloride/methanol = 9:1)

$C_{22}H_{23}N_3O_2$

30 Mass spectrum: $m/z = 361 (m^+)$

(27) 3-Z-[1-(4-amino-cyclohexyl-amino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-amino-cyclohexylamine
 $C_{22}H_{24}N_4O_2$

5 Mass spectrum: $m/z = 376 (m^+)$

(28) 3-Z-[1-(N-methyl-piperidine-4-yl-amino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 4-amino-1-methyl-piperidine
10

R_f value: 0.15 (silica gel, methylene chloride/methanol = 4:1)

$C_{22}H_{24}N_4O_2$

Mass spectrum: $m/z = 376 (m^+)$

15 (29) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II(2) and 4-(piperidin-1-yl-methyl)-aniline

R_f value: 0.30 (silica gel, methylene chloride/methanol = 4:1)

20 $C_{23}H_{26}N_4O_2$

Mass spectrum: $m/z = 390 (m^+)$

(30) 3-Z-[1-(3-dimethylaminomethyl-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

25 Prepared from the resin obtained according to Example II(2) and 3-dimethylaminomethyl-aniline

R_f value: 0.51 (silica gel, methylene chloride/methanol = 4:1)

$C_{20}H_{22}N_4O_2$

Mass spectrum: $m/z = 351 (m+H^+)$

30

(31) 3-Z-[1-(4-(N-methyl-N-benzyl-aminomethyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II(2) and 4-(N-methyl-N-benzyl-aminomethyl)-aniline

5 R_f value: 0.73 (silica gel, methylene chloride/methanol = 4:1)

C₂₆H₂₆N₄O₂

Mass spectrum: m/z = 426 (m⁺)

10 (32) 3-Z-[1-(4-(N-methylsulphonyl-N-(2-dimethylamino-ethyl)-amino)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II(2) and 4-(N-methylsulphonyl-N-(2-dimethylamino-ethyl)-amino)-aniline

C₂₂H₂₇N₅O₄S

Mass spectrum: m/z = 458 (m+H⁺)

15

(33) 3-Z-[1-(4-chloro-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(2) and 4-chloro-aniline

R_f value: 0.10 (silica gel, methylene chloride/methanol = 9:1)

C₁₇H₁₄ClN₃O₂

20 Mass spectrum: m/z = 327/329 (m⁺)

(34) 3-Z-[1-(3-chloro-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(2) and 3-chloro-aniline

R_f value: 0.11 (silica gel, methylene chloride/methanol = 9:1)

25 C₁₇H₁₄ClN₃O₂

Mass spectrum: m/z = 327/329 (m⁺)

(35) 3-Z-[1-(4-methoxycarbonyl-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(2) and methyl 4-amino-benzoate

30 R_f value: 0.11 (silica gel, methylene chloride/methanol = 9:1)

C₁₉H₁₇N₃O₄

Mass spectrum: $m/z = 351$ (m^+)

(36) 3-Z-[1-(4-carboxy-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(2) and 4-amino-benzoic acid

5 $C_{18}H_{15}N_3O_4$

Mass spectrum: $m/z = 336$ ($m-H^+$)

(37) 3-Z-[1-(4-methyl-3-nitro-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(2) and 4-methyl-3-nitro-aniline

10 R_f value: 0.82 (silica gel, methylene chloride/methanol = 4:1)

$C_{18}H_{16}N_4O_4$

Mass spectrum: $m/z = 352$ (m^+)

15 (38) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II(4) and 4-(piperidin-1-yl-methyl)-aniline

R_f value: 0.37 (silica gel, methylene chloride/methanol = 4:1)

$C_{25}H_{30}N_4O_2$

20 Mass spectrum: $m/z = 418$ (m^+)

(39) 3-Z-[1-(3-dimethylaminomethyl-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

25 Prepared from the resin obtained according to Example II(4) and 3-dimethylaminomethyl-aniline

R_f value: 0.42 (silica gel, methylene chloride/methanol = 4:1)

$C_{22}H_{26}N_4O_2$

Mass spectrum: $m/z = 378$ (m^+)

30 (40) 3-Z-[1-(4-(N-methyl-N-benzyl-aminomethyl)-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II(4) and 4-(N-methyl-N-benzyl-aminomethyl)-aniline

R_f value: 0.81 (silica gel, methylene chloride/methanol = 4:1)

C₂₈H₃₀N₄O₂

5 Mass spectrum: m/z = 454 (m⁺)

(41) 3-Z-[1-(4-(N-methylsulphonyl-N-(2-dimethylamino-ethyl)-amino)-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II(4) and 4-(N-methylsulphonyl-N-(2-dimethylamino-ethyl)-amino)-aniline

10

R_f value: 0.59 (silica gel, methylene chloride/methanol = 4:1)

C₂₄H₃₁N₅O₄S

Mass spectrum: m/z = 486 (m+H⁺)

15 (42) 3-Z-[1-(4-chloro-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(4) and 4-chloro-aniline

R_f value: 0.17 (silica gel, methylene chloride/methanol = 9:1)

C₁₉H₁₈ClN₃O₂

Mass spectrum: m/z = 355/357 (m⁺)

20

(43) 3-Z-[1-(3-chloro-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(4) and 3-chloro-aniline

R_f value: 0.12 (silica gel, methylene chloride/methanol = 9:1)

C₁₉H₁₈ClN₃O₂

25 Mass spectrum: m/z = 355/357 (m⁺)

(44) 3-Z-[1-(4-methoxycarbonyl-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(4) and methyl 4-amino-benzoate

R_f value: 0.8 (silica gel, methylene chloride/methanol = 4:1)

30 C₂₁H₂₁N₃O₄

Mass spectrum: m/z = 379 (m⁺)

(45) 3-Z-[1-(4-carboxy-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(4) and 4-amino-benzoic acid

$C_{20}H_{19}N_3O_4$

5 Mass spectrum: $m/z = 364$ ($m-H^+$)

(46) 3-Z-[1-(4-methyl-3-nitro-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

Prepared from the resin obtained according to Example II(4) and 4-methyl-3-nitro-aniline

R_f value: 0.86 (silica gel, methylene chloride/methanol = 4:1)

10 $C_{20}H_{20}N_4O_4$

Mass spectrum: $m/z = 380$ (m^+)

Example 2

15 3-Z-[1-(3-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

2.0 g of resin obtained according to Example II are reacted analogously to Example 1 with 2.0 g of 3-aminobenzyl alcohol in 20 ml of dimethylformamide for 22 hours at 70° C. Then
 20 the solvent is suction filtered and the resin is washed several times with dimethylformamide and methylene chloride. Then 200 mg of the moist charged resin are suspended in 2 ml of methylene chloride and left to stand with 0.2 ml of methanesulphonic acid chloride and 0.1 ml of triethylamine for 2 hours at room temperature. Then the resin is washed several times with methylene chloride, suspended in 2 ml of methylene chloride
 25 and combined with 0.2 ml of piperidine. After 1 hour the resin is washed with methylene chloride and dimethylformamide and then treated with trifluoroacetic acid analogously to Example 1.

Yield: 15 mg

R_f value: 0.30 (silica gel, methylene chloride/methanol = 4:1)

30 $C_{28}H_{28}N_4O_2$

Mass spectrum: $m/z = 452$ (m^+)

The following compounds are prepared analogously to Example 2:

- 5 (1) 3-Z-[1-(3-(diethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and diethylamine

R_f value: 0.80 (silica gel, methylene chloride/methanol = 4:1)

C₂₇H₂₈N₄O₂

Mass spectrum: m/z = 440 (m⁺)

10

- (2) 3-Z-[1-(3-(benzylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and benzylamine

R_f value: 0.80 (silica gel, methylene chloride/methanol = 4:1)

- 15 C₃₀H₂₆N₄O₂

Mass spectrum: m/z = 474 (m⁺)

- (3) 3-Z-[1-(3-(N-methyl-N-benzyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

- 20 Prepared from the resin obtained according to Example II and N-methyl-benzylamine

R_f value: 0.80 (silica gel, methylene chloride/methanol = 4:1)

C₃₁H₂₈N₄O₂

Mass spectrum: m/z = 488 (m⁺)

- 25 (4) 3-Z-[1-(3-(butylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and butylamine

R_f value: 0.40 (silica gel, methylene chloride/methanol = 4:1)

C₂₇H₂₈N₄O₂

- 30 Mass spectrum: m/z = 440 (m⁺)

(5) 3-Z-[1-(3-(aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and ammonia

$C_{23}H_{20}N_4O_2$

5 Mass spectrum: $m/z = 385 (m+H^+)$

(6) 3-Z-[1-(3-(N-(3-dimethylaminopropyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 1-dimethylamino-3-

10 methylaminopropane

R_f value: 0.67 (silica gel, methylene chloride/methanol = 4:1)

$C_{29}H_{33}N_5O_2$

Mass spectrum: $m/z = 484 (m+H^+)$

15 (7) 3-Z-[1-(3-(N-(2-dimethylaminoethyl)-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

Prepared from the resin obtained according to Example II and 1-dimethylamino-2-methylaminoethane

R_f value: 0.40 (silica gel, methylene chloride/methanol = 4:1)

20 $C_{28}H_{31}N_5O_2$

Mass spectrum: $m/z = 470 (m+H^+)$

Example 3

3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5

1.5 g of 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 1.1 g of 4-(piperidin-1-yl-methyl)-aniline are dissolved in 15 ml of dimethylformamide and stirred for 45 minutes at 100°C. After cooling 5.0 ml of piperidine are added and the mixture is stirred for another 3 hours at room temperature. The solvent is removed and the residue purified over an aluminium oxide column (activity: 2-3) with methylene chloride/ethanol (100:3) as eluant.

10

Yield: 1.1 g (58% of theory),

R_f value: 0.5 (aluminium oxide, methylene chloride/ethanol = 100:3)

C₃₀H₃₁N₃O₃

15

Mass spectrum: m/z = 481 [M⁺]

The following compounds are prepared analogously to Example 3:

(1) 3-Z-[1-(4-bromo-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone

20

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-bromoaniline

R_f value: 0.4 (silica gel, toluene/ethyl acetate = 5:1)

C₂₄H₁₉BrN₂O₃

Mass spectrum: m/z = 462/464 [M⁺]

25

(2) 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 3-(dimethylaminomethyl)-aniline

30

R_f value: 0.5 (aluminium oxide, methylene chloride/ethanol = 30:1)

C₂₇H₂₇N₃O₃

ESI mass spectrum: $m/z = 442$ $[M+H^+]$

(3) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

- 5 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(dimethylaminomethyl)-aniline

R_f value: 0.7 (aluminium oxide, ethyl acetate/ethanol = 20:1)

$C_{27}H_{27}N_3O_3$

ESI mass spectrum: $m/z = 442$ $[M+H^+]$

10

(4) 3-Z-[1-(4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-aniline

- 15 R_f value: 0.6 (silica gel, methylene chloride/ethanol = 5:1)

$C_{32}H_{35}N_3O_3$

Mass spectrum: $m/z = 509$ $[M^+]$

(5) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(2-dimethylamino-ethyl)-aniline

R_f value: 0.2 (silica gel, methylene chloride/ethanol = 5:1)

$C_{28}H_{29}N_3O_3$


- 25 Mass spectrum: $m/z = 455$ $[M^+]$

(6) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-aniline

R_f value: 0.4 (aluminium oxide, methylene chloride/ethanol = 20:1)




Mass spectrum: $m/z = 512 [M^+]$

- 5 (7) 3-Z-[1-(4-tert.butyloxycarbonyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-tert.butyloxycarbonyl-aniline

R_f value: 0.4 (aluminium oxide, methylene chloride/ethanol = 40:1)




- 10 Mass spectrum: $m/z = 484 [M^+]$

- (8) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-aniline

R_f value: 0.2 (aluminium oxide, methylene chloride/ethanol = 40:1)



Mass spectrum: $m/z = 526 [M^+]$

- 20 (9) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and N-(2-dimethylamino-ethyl)-N-methylsulphonyl-p-phenylenediamine

R_f value: 0.3 (aluminium oxide, methylene chloride/ethanol = 40:1)

- 25 $C_{29}H_{32}N_4O_5S$

Mass spectrum: $m/z = 548 [M^+]$

- (10) 3-Z-[1-(4-(4-methyl-piperazin-1-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

- 30 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(4-methyl-piperazin-1-yl)-aniline

R_f value: 0.3 (aluminium oxide, ethyl acetate)

C₂₉H₃₀N₄O₃

ESI mass spectrum: m/z = 483 [M+H⁺]

- 5 (11) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-methyl-amino)-aniline

R_f value: 0.5 (aluminium oxide, methylene chloride/ethanol = 20:1)

- 10 C₂₉H₃₂N₄O₃

ESI mass spectrum: m/z = 485 [M+H⁺]

- (12) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(N-(3-dimethylamino-propyl)-N-methyl-amino)-aniline

R_f value: 0.5 (aluminium oxide, ethyl acetate)

C₃₀H₃₄N₄O₃

ESI mass spectrum: m/z = 499 [M+H⁺]

- 20

- (13) 3-Z-[1-(4-(N-methyl-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-amino-N-methyl-acetanilide

- 25 R_f value: 0.3 (silica gel, methylene chloride/ethanol = 15:1)

C₂₇H₂₅N₃O₄

Mass spectrum: m/z = 455 [M⁺]

- 30 (14) 3-Z-[1-(4-(N-methyl-methylsulphonylamino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and N-(4-aminophenyl)-N-methyl-methanesulphonamide

R_f value: 0.8 (aluminium oxide, ethyl acetate)

C₂₆H₂₅N₃O₅S

5 Mass spectrum: m/z = 491 [M⁺]

(15) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and N-(3-dimethylamino-propyl)-N-methylsulphonyl-p-phenylenediamine

R_f value: 0.6 (silica gel, methylene chloride/ethanol/ammonia = 5:2:0.01)

C₃₀H₃₄N₄O₅S

ESI mass spectrum: m/z = 563 [M+H⁺]

15 (16) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl)-amino)-aniline

R_f value: 0.6 (silica gel, methylene chloride/ethanol = 10:1)

20 C₂₉H₃₀N₄O₆S

ESI mass spectrum: m/z = 561 [M-H]

(17) 3-Z-[1-(4-(imidazol-4-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(imidazol-4-yl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/ethanol/ammonia = 10:1:0.01)

C₂₇H₂₂N₄O₃

Mass spectrum: m/z = 450 [M⁺]

30

(18) 3-Z-[1-(4-(tetrazol-5-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(tetrazol-5-yl)-aniline

5 R_f value: 0.5 (silica gel, methylene chloride/ethanol = 5:1)

$C_{25}H_{20}N_6O_3$

ESI mass spectrum: $m/z = 451$ $[M-H]^-$

(19) 3-Z-[1-(4-(N-benzyl-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(N-benzyl-N-methyl-aminomethyl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/ethanol = 10:1)

$C_{33}H_{31}N_3O_3$

15 ESI mass spectrum: $m/z = 516$ $[M-H]^-$

(20) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-propionyl-amino)-aniline)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-[N-(2-dimethylamino-ethyl)-N-propionyl-amino]-aniline

20

R_f value: 0.2 (silica gel, methylene chloride/ethanol = 5:1)

$C_{31}H_{34}N_4O_4$

ESI mass spectrum: $m/z = 525$ $[M-H]^-$

(21) 3-Z-[1-(4-(pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and 4-(pyrrolidin-1-yl-methyl)-aniline

R_f value: 0.1 (silica gel, methylene chloride/ethanol = 5:1)

30 $C_{29}H_{29}N_3O_3$

ESI mass spectrum: $m/z = 466$ $[M-H]^-$

(22) 3-Z-[1-(4-(N-methyl-N-phenethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone
5 and 4-(N-phenethyl-N-methyl-aminomethyl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/ethanol = 10:1)

C₃₄H₃₃N₃O₃

ESI mass spectrum: m/z = 530 [M-H]

10 (23) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone
and N-dimethylaminomethylcarbonyl-N-methyl-p-phenylenediamine

R_f value: 0.1 (silica gel, methylene chloride/ethanol = 10:1)

15 C₂₉H₃₀N₄O₄

ESI mass spectrum: m/z = 497 [M-H]

(24) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-ethylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone
and N-(2-dimethylamino-ethyl)-N-ethylsulphonyl-p-phenylenediamine

R_f value: 0.6 (silica gel, methylene chloride/ethanol = 5:1)

C₃₀H₃₄N₄O₅S

ESI mass spectrum: m/z = 561 [M-H]

25

(25) 3-Z-[1-(4-(N-tert.butoxycarbonyl-N-ethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone
and 4-(N-tert.butoxycarbonyl-N-ethyl-aminomethyl)-aniline

30 R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

C₃₂H₃₅N₃O₅

ESI mass spectrum: $m/z = 540$ $[M-H^-]$

(26) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

- 5 Prepared from 1-acetyl-3-(1-ethoxy-1-ethyl-methylene)-6-ethoxycarbonyl-2-indolinone and 4-(piperidin-1-yl-methyl)-aniline

R_f value: 0.9 (silica gel, methylene chloride/ethanol = 5:1)

$C_{26}H_{31}N_3O_3$

ESI mass spectrum: $m/z = 432$ $[M-H^-]$

10

(27) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-ethyl-methylene)-6-ethoxycarbonyl-2-indolinone and N-(2-dimethylamino-ethyl)-N-methylsulphonyl-p-phenylenediamine

- 15 R_f value: 0.3 (silica gel, methylene chloride/ethanol = 5:1)

$C_{25}H_{32}N_4O_5S$

ESI mass spectrum: $m/z = 499$ $[M-H^-]$

(28) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(dimethylaminomethyl)-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol = 5:1)


$C_{26}H_{25}N_3O_3$

- 25 ESI mass spectrum: $m/z = 428$ $[M+H^+]$

(29) 3-Z-[1-(4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 30 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-aniline

R_f value: 0.5 (RP 8, methanol/five percent saline solution = 4:1)




ESI mass spectrum: $m/z = 496 [M+H^+]$

- (30) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(2-dimethylamino-ethyl)-N-methylsulphonyl-p-phenylenediamine

R_f value: 0.6 (silica gel, methylene chloride/methanol = 5:1)




- ESI mass spectrum: $m/z = 533 [M-H^-]$

- (31) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(3-dimethylamino-propyl)-N-methylsulphonyl-p-phenylenediamine

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 30:1)



ESI mass spectrum: $m/z = 547 [M-H^-]$

- (32) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-dimethylaminocarbonyl-methyl-N-methylsulphonyl)-amino)-aniline

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 20:1)

- $C_{28}H_{28}N_4O_6S$

ESI mass spectrum: $m/z = 547 [M-H^-]$

- (33) 3-Z-[1-(4-(N-acetyl-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-acetyl-N-dimethylaminocarbonylmethyl)-amino)-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol = 10:1)

C₂₉H₂₈N₄O₅

ESI mass spectrum: m/z = 511 [M-H]⁻

- 5 (34) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-dimethylaminocarbonyl-methyl)-amino)-aniline

R_f value: 0.6 (aluminium oxide, methylene chloride/methanol = 30:1)

- 10 C₂₇H₂₆N₄O₄

ESI mass spectrum: m/z = 469 [M-H]⁻

(35) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(3-dimethylamino-propyl)-N-acetyl-p-phenylenediamine

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 20:1)

C₃₀H₃₂N₄O₄

ESI mass spectrum: m/z = 511 [M-H]⁻

- 20

(36) 3-Z-[1-(4-(N-methylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-methylaminocarbonylmethyl-N-methylsulphonyl-p-phenylenediamine

- 25 R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

C₂₇H₂₆N₄O₆S

ESI mass spectrum: m/z = 533 [M-H]⁻

- 30 (37) 3-Z-[1-(4-((imidazolidin-2,4-dione-5-ylidene)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-((imidazolidin-2,4-dione-5-ylidene)-methyl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/methanol = 10:1)

C₂₇H₂₀N₄O₅

5 ESI mass spectrum: m/z = 479 [M-H⁻]

(38) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
10 and N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-p-phenylenediamine

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 20:1)

C₂₉H₃₀N₄O₄

ESI mass spectrum: m/z = 497 [M-H⁻]

15 (39) 3-Z-[1-(4-(N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(N-tert.butoxycarbonyl-aminomethyl)-aniline

R_f value: 0.3 (aluminium oxide, methylene chloride/methanol = 20:1)

20 C₂₉H₂₉N₃O₅

ESI mass spectrum: m/z = 498 [M-H⁻]

(40) 3-Z-[1-(4-(2-oxo-pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(2-oxo-pyrrolidin-1-yl-methyl)-aniline

R_f value: 0.3 (silica gel, methylene chloride/methanol = 20:1)

C₂₈H₂₅N₃O₄

ESI mass spectrum: m/z = 466 [M-H⁻]

30

(41) 3-Z-[1-(4-(N-aminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-aminocarbonylmethyl-N-methylsulphonyl-p-phenylenediamine

5 R_f value: 0.7 (silica gel, methylene chloride/methanol = 5:1)

C₂₆H₂₄N₄O₆S

ESI mass spectrum: m/z = 519 [M-H]⁻

(42) 3-Z-[1-(4-(thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(thiomorpholin-4-yl-methyl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/methanol = 15:1)

C₂₈H₂₇N₃O₃S

15 ESI mass spectrum: m/z = 484 [M-H]⁻

(43) 3-Z-[1-(4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

C₂₈H₂₇N₃O₅S

ESI mass spectrum: m/z = 516 [M-H]⁻

25 (44) 3-Z-[1-(4-(N-cyanomethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-cyanomethyl-N-methyl-sulphonyl-p-phenylenediamine

R_f value: 0.6 (silica gel, methylene chloride/methanol = 10:1)

30 C₂₆H₂₂N₄O₅S

ESI mass spectrum: m/z = 501 [M-H]⁻

(45) 3-Z-[1-(4-(N-tert.butoxycarbonyl-ethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
5 and 4-(N-ethyl-N-tert.butoxycarbonyl-aminomethyl)-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol = 10:1)

C₃₁H₃₃N₃O₅

ESI mass spectrum: m/z = 526 [M-H]

10 (46) 3-Z-[1-(4-(N-benzyl-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(N-benzyl-N-methyl-aminomethyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

15 C₃₂H₂₉N₃O₃

ESI mass spectrum: m/z = 502 [M-H]

(47) 3-Z-[1-(4-(1-oxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(1-oxo-thiomorpholin-4-yl-methyl)-aniline

R_f value: 0.7 (silica gel, methylene chloride/methanol = 10:1)

C₂₈H₂₇N₃O₄S

ESI mass spectrum: m/z = 500 [M-H]

25

(48) 3-Z-[1-(4-(2-(imidazol-4-yl)-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(2-(imidazol-4-yl)-ethyl)-aniline

30 R_f value: 0.4 (silica gel, methylene chloride/methanol = 5:1)

C₂₈H₂₄N₄O₃

ESI mass spectrum: $m/z = 463$ [M-H]

(49) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 5 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(morpholin-4-yl-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

$C_{28}H_{27}N_3O_4$

ESI mass spectrum: $m/z = 468$ [M-H]

10

(50) 3-Z-[1-(4-((4-methyl-piperazin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((4-methyl-piperazin-1-yl)-methyl)-aniline

- 15 R_f value: 0.4 (silica gel, methylene chloride/methanol 5:1)

$C_{29}H_{30}N_4O_3$

ESI mass spectrum: $m/z = 481$ [M-H]

(51) 3-Z-[1-(4-((2-(N-benzyl-N-methyl-amino)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-(N-benzyl-N-methyl-amino)-ethyl)-N-methylsulphonyl-amino)-aniline

R_f value: 0.7 (silica gel, methylene chloride/methanol = 10:1)

$C_{34}H_{34}N_4O_5S$


- 25 ESI mass spectrum: $m/z = 609$ [M-H]

(52) 3-Z-[1-(4-cyclohexylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-cyclohexyl-p-phenylenediamine

R_f value: 0.8 (silica gel, methylene chloride/methanol = 10:1)




ESI mass spectrum: $m/z = 451$ $[M-H]^-$

- 5 (53) 3-Z-[1-(4-(pyridin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(pyridin-4-yl-methyl)-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)




- 10 ESI mass spectrum: $m/z = 460$ $[M-H]^-$

- (54) 3-Z-[1-(4-(imidazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(imidazol-1-yl-methyl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/methanol/ammonia = 10:1:0.01)



ESI mass spectrum: $m/z = 449$ $[M-H]^-$

- 20 (55) 3-Z-[1-(4-(imidazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(imidazol-1-yl-methyl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/methanol/ammonia = 10:1:0.01)

- 25 $C_{27}H_{22}N_4O_3$

ESI mass spectrum: $m/z = 449$ $[M-H]^-$

- (56) 3-Z-[1-(N-methyl-piperidine-4-yl-amino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 30 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-amino-1-methyl-piperidine

R_f value: 0.3 (silica gel, methylene chloride/methanol = 5:1)

C₂₃H₂₅N₃O₃

ESI mass spectrum: m/z = 390 [M-H⁻]

- 5 (57) 3-Z-[1-(4-(imidazol-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(imidazol-4-yl-methyl)-aniline

R_f value: 0.2 (silica gel, methylene chloride/methanol = 5:1)

- 10 C₂₇H₂₂N₄O₃

ESI mass spectrum: m/z = 449 [M-H⁻]

(58) 3-Z-[1-(4-((4-hydroxy-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((4-hydroxy-piperidin-1-yl)-methyl)-aniline

R_f value: 0.1 (silica gel, methylene chloride/methanol = 10:1)

C₂₉H₂₉N₃O₃

ESI mass spectrum: m/z = 482 [M-H⁻]

- 20

(59) 3-Z-[1-(4-((4-methoxy-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((4-methoxy-piperidin-1-yl)-methyl)-aniline

- 25 R_f value: 0.4 (silica gel, methylene chloride/methanol = 10:1)

C₃₀H₃₁N₃O₄

ESI mass spectrum: m/z = 496 [M-H⁻]

(60) 3-Z-[1-(4-benzyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 30 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-benzyl-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol = 10:1)

C₃₀H₂₄N₂O₃

Melting point: 224°C

- 5 (61) 3-Z-[1-(4-(N-(3-trifluoroacetyl-amino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(3-trifluoroacetyl-amino-propyl)-N-methylsulphonyl-p-phenylenediamine

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 20:1)

- 10 C₂₉H₂₇F₃N₄O₆S

ESI mass spectrum: m/z = 615 [M-H⁻]

- (62) 3-Z-[1-(4-tert.butoxycarbonylmethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-ethoxycarbonyl-2-indolinone and tert.butyl 4-aminophenylacetate

R_f value: 0.5 (aluminium oxide, ethyl acetate)

C₃₀H₃₀N₂O₅

ESI mass spectrum: m/z = 497 [M-H⁻]

- 20

- (63) 3-Z-[1-(4-tert.butoxycarbonyl-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-ethylmethylen)-6-ethoxycarbonyl-2-indolinone and 4-tert.butoxycarbonyl-aniline

- 25 R_f value: 0.4 (aluminium oxide, methylene chloride/ethanol = 20:1)

C₂₅H₂₈N₂O₅

ESI mass spectrum: m/z = 435 [M-H⁻]

- 30 (64) 3-Z-[1-(4-(4-tert.butoxycarbonyl-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(4-tert.butoxycarbonyl-piperazin-1-yl-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

C₃₃H₃₆N₄O₅

5 ESI mass spectrum: m/z = 567 [M-H⁻]

(65) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
10 and 4-(1-methyl-imidazol-2-yl)-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol = 5:1)

C₂₇H₂₂N₄O₃

ESI mass spectrum: m/z = 449 [M-H⁻]

15 (66) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-nitro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 6-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-amino-nitrobenzene

R_f value: 0.6 (silica gel, methylene chloride/methanol = 5:1)

20 C₂₈H₂₉N₅O₇S

ESI mass spectrum: m/z = 578 [M-H⁻]

(67) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-amino-aniline

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 20:1)

C₂₈H₃₁N₅O₅S

ESI mass spectrum: m/z = 548 [M-H⁻]

30

(68) 3-Z-[1-(4-((3-(N-benzyl-N-methyl-amino)-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(3-(N-benzyl-N-methyl-amino)-propyl)-N-methylsulphonyl-amino)-aniline

5 R_f value: 0.6 (silica gel, methylene chloride/methanol = 10:1)

C₃₅H₃₆N₄O₅S

ESI mass spectrum: m/z = 623 [M-H⁻]

(69) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-chloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-chloro-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

C₂₈H₂₉ClN₄O₅S

15 ESI mass spectrum: m/z = 567/569 [M-H⁻]

(70) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-dimethylaminomethylcarbonyl-N-methyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₂₈H₂₈N₄O₄

ESI mass spectrum: m/z = 483 [M-H⁻]

(71) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

30 C₂₉H₃₀N₄O₄

ESI mass spectrum: m/z = 497 [M-H⁻]

(72) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-propionyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
5 and 4-(N-(2-dimethylamino-ethyl)-N-propionyl-amino)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₃₀H₃₂N₄O₄

ESI mass spectrum: m/z = 511 [M-H⁻]

10 (73) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-butyryl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(N-(2-dimethylamino-ethyl)-N-butyryl-amino)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

15 C₃₁H₃₄N₄O₄

ESI mass spectrum: m/z = 525 [M-H⁻]

(74) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-isobutyryl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(N-(2-dimethylamino-ethyl)-N-isobutyryl-amino)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₃₁H₃₄N₄O₄

ESI mass spectrum: m/z = 525 [M-H⁻]

25

(75) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-benzoyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(N-(2-dimethylamino-ethyl)-N-benzoyl-amino)-aniline

30 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₃₄H₃₂N₄O₄

ESI mass spectrum: $m/z = 559$ $[M-H]^-$

(76) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 5 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-amino-aniline

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 20:1)

$C_{29}H_{31}N_5O_4$

ESI mass spectrum: $m/z = 512$ $[M-H]^-$

10

(77) 3-Z-[1-(4-(4-hydroxymethyl-piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(4-hydroxymethyl-piperidin-1-yl-methyl-amino)-aniline

- 15 R_f value: 0.3 (silica gel, methylene chloride/methanol = 5:1)

$C_{30}H_{31}N_3O_4$

ESI mass spectrum: $m/z = 496$ $[M-H]^-$

(78) 3-Z-[1-(4-(2-(4-hydroxy-piperidin-1-yl)-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(2-(4-hydroxy-piperidin-1-yl)-ethyl-amino)-aniline

R_f value: 0.3 (silica gel, methylene chloride/methanol = 5:1)

$C_{30}H_{31}N_3O_4$

- 25 ESI mass spectrum: $m/z = 496$ $[M-H]^-$

(79) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-propylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 30 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(2-dimethylamino-ethyl)-N-propylsulphonyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{30}H_{34}N_4O_5S$

ESI mass spectrum: $m/z = 561 [M-H^-]$

- 5 (80) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-butylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(2-dimethylamino-ethyl)-N-butylsulphonyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{31}H_{36}N_4O_5S$

- 10 ESI mass spectrum: $m/z = 575 [M-H^-]$

(81) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-phenylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(2-dimethylamino-ethyl)-N-phenylsulphonyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{33}H_{32}N_4O_5S$

ESI mass spectrum: $m/z = 595 [M-H^-]$

- 20 (82) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-benzylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(2-dimethylamino-ethyl)-N-benzylsulphonyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

- 25 $C_{34}H_{34}N_4O_5S$

ESI mass spectrum: $m/z = 609 [M-H^-]$

(83) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-ethylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 30 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(2-dimethylamino-ethyl)-N-ethylsulphonyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₂₉H₃₂N₄O₅S

ESI mass spectrum: m/z = 547 [M-H⁻]

- 5 (84) 3-Z-[1-(4-((imidazolidin-2,4-dion-5-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((imidazolidin-2,4-dion-5-yl)-methyl)-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol = 5:1)

- 10 C₂₇H₂₂N₄O₅

ESI mass spectrum: m/z = 481 [M-H⁻]

- (85) 3-Z-[1-(4-((3-hydroxy-pyrrolidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((3-hydroxy-pyrrolidin-1-yl)-methyl)-aniline

R_f value: 0.1 (silica gel, methylene chloride/methanol = 10:1)

C₂₈H₂₇N₃O₄

ESI mass spectrum: m/z = 468 [M-H⁻]

- 20

- (86) 3-Z-[1-(4-(cyclohexyl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(cyclohexyl-methyl)-aniline (Eur. J. Med. Chem. Chim. Ther. 1992, 27, 537-544)

- 25 R_f value: 0.6 (silica gel, methylene chloride/methanol = 10:1)

C₃₀H₃₀N₂O₃

ESI mass spectrum: m/z = 465 [M-H⁻]

- (87) 3-Z-[1-(4-(cyclohexyl-carbonyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 30

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(cyclohexyl-carbonyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

C₃₀H₂₈N₂O₄

5 ESI mass spectrum: m/z = 479 [M-H⁻]

(88) 3-Z-[1-(4-diethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(diethylamino-methyl)-aniline

10 R_f value: 0.4 (silica gel, methylene chloride/methanol = 10:1)

C₂₈H₂₉N₃O₃

ESI mass spectrum: m/z = 454 [M-H⁻]

15 (89) 3-Z-[1-(4-(N-(n-hexyl)-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(n-hexyl)-N-methyl-aminomethyl)-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol = 10:1)

20 C₃₁H₃₅N₃O₃

ESI mass spectrum: m/z = 496 [M-H⁻]

(90) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(furan-2-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-(furan-2-carbonyl)-amino)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₃₂H₃₀N₄O₅

ESI mass spectrum: m/z = 549 [M-H⁻]

30

(91) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(2-methoxy-benzoyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-(2-methoxy-benzoyl)-amino)-aniline

5 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{35}H_{34}N_4O_5$

ESI mass spectrum: $m/z = 589 [M-H^-]$

(92) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(pyridine-3-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-(pyridine-3-carbonyl)-amino)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{33}H_{31}N_5O_4$

15 ESI mass spectrum: $m/z = 560 [M-H^-]$

(93) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(phenyl-acetyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-(phenyl-acetyl)-amino)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{35}H_{34}N_4O_4$

ESI mass spectrum: $m/z = 573 [M-H^-]$

25 (94) 3-Z-[1-(4-(N-ethyl-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-ethyl-N-methyl-aminomethyl)-aniline

R_f value: 0.3 (silica gel, methylene chloride/methanol = 10:1)

30 $C_{27}H_{27}N_3O_3$

ESI mass spectrum: $m/z = 440 [M-H^-]$

(95) 3-Z-[1-(4-(imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
5 and 4-(imidazol-2-yl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

C₂₆H₂₀N₄O₃

ESI mass spectrum: m/z = 435 [M-H⁻]

10 (96) 3-Z-[1-(4-(1-ethyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(1-ethyl-imidazol-2-yl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/methanol = 10:1)

15 C₂₈H₂₄N₄O₃

ESI mass spectrum: m/z = 463 [M-H⁻]

(97) 3-Z-[1-(4-(1-benzyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(1-benzyl-imidazol-2-yl)-aniline

R_f value: 0.3 (silica gel, methylene chloride/methanol = 20:1)

C₃₃H₂₆N₄O₃

ESI mass spectrum: m/z = 525 [M-H⁻]

25

(98) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-isopropylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and N-(2-dimethylamino-ethyl)-N-isopropylsulphonyl-p-phenylenediamine

30 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₃₀H₃₄N₄O₅S

ESI mass spectrum: $m/z = 561$ $[M-H]^-$

(99) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 5 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(piperidin-1-yl-methylcarbonyl)-N-methyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{31}H_{32}N_4O_4$

ESI mass spectrum: $m/z = 523$ $[M-H]^-$

10

(100) 3-Z-[1-(4-(N-(morpholin-4-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(morpholin-4-yl-methylcarbonyl)-N-methyl-p-phenylenediamine

- 15 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{30}H_{30}N_4O_5$

ESI mass spectrum: $m/z = 525$ $[M-H]^-$

(101) 3-Z-[1-(4-(N-((4-benzyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((4-benzyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{37}H_{37}N_5O_4$


- 25 ESI mass spectrum: $m/z = 614$ $[M-H]^-$

(102) 3-Z-[1-(4-(N-(pyrrolidin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(pyrrolidin-1-yl-methylcarbonyl)-N-methyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)




ESI mass spectrum: $m/z = 509 [M-H^-]$

- 5 (103) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-bromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-bromo-aniline

R_f value: 0.6 (silica gel, methylene chloride/methanol = 5:1)




- 10 ESI mass spectrum: $m/z = 575/577 [M-H^-]$

- (104) 3-Z-[1-(4-(5-methyl-imidazol-4-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(5-methyl-imidazol-4-yl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol/ammonia = 10:1:0.01)



ESI mass spectrum: $m/z = 449 [M-H^-]$

- 20 (105) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((2-dimethylamino-ethyl)-carbonyl)-N-isopropyl-p-phenylenediamine

R_f value: 0.1 (silica gel, methylene chloride/methanol = 10:1)

- 25 $C_{31}H_{34}N_4O_4$

ESI mass spectrum: $m/z = 525 [M-H^-]$

- (106) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 30 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((2-dimethylamino-ethyl)-carbonyl)-N-benzyl-p-phenylenediamine

R_f value: 0.1 (silica gel, methylene chloride/methanol = 10:1)

C₃₁H₃₄N₄O₄

ESI mass spectrum: m/z = 525 [M-H⁻]

- 5 (107) 3-Z-[1-(4-(N-butyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(N-butyl-N-tert.butoxycarbonyl-aminomethyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

- 10 C₃₃H₃₇N₃O₅

ESI mass spectrum: m/z = 554 [M-H⁻]

(108) 3-Z-[1-(4-(N-((N-aminocarbonylmethyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(N-aminocarbonylmethyl-N-methyl-amino)-methylcarbonyl)-N-methyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₂₉H₂₉N₅O₅

- 20 ESI mass spectrum: m/z = 526 [M-H⁻]

(109) 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-methyl-p-phenylenediamine

- 25 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₃₄H₃₂N₄O₄

ESI mass spectrum: m/z = 559 [M-H⁻]

- 30 (110) 3-Z-[1-(4-(N-(di-(2-methoxyethyl)-amino-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(di-(2-methoxyethyl)-amino-methylcarbonyl)-N-methyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₃₂H₃₆N₄O₆

5 ESI mass spectrum: m/z = 571 [M-H⁻]

(111) 3-Z-[1-(4-(N-((2-(4-tert.butoxycarbonyl-piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((2-(4-tert.butoxycarbonyl-piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-p-phenylenediamine

R_f value: 0.8 (silica gel, methylene chloride/methanol = 5:1)

C₃₆H₄₁N₅O₆

ESI mass spectrum: m/z = 638 [M-H⁻]

15

(112) 3-Z-[1-(4-(N-((2-(piperidin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((2-(piperidin-1-yl)-ethyl)-carbonyl)-N-methyl-p-phenylenediamine

20 R_f value: 0.5 (silica gel, methylene chloride/methanol = 5:1)

C₃₂H₃₄N₄O₄

ESI mass spectrum: m/z = 537 [M-H⁻]

(113) 3-Z-[1-(4-(N-((2-(N-benzyl-N-methyl-amino)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((2-(N-benzyl-N-methyl-amino)-ethyl)-carbonyl)-N-methyl-p-phenylenediamine

R_f value: 0.4 (silica gel, methylene chloride/methanol = 10:1)

C₃₅H₃₄N₄O₄

30 ESI mass spectrum: m/z = 573 [M-H⁻]

(114) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(dimethylaminomethylcarbonyl)-N-isopropyl-p-phenylenediamine

5 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{30}H_{32}N_4O_4$

ESI mass spectrum: $m/z = 511 [M-H^-]$

10 (115) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(piperidin-1-yl-methylcarbonyl)-N-isopropyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{33}H_{36}N_4O_4$

15 ESI mass spectrum: $m/z = 551 [M-H^-]$

(116) 3-Z-[1-(4-(N-((4-tert.butoxycarbonyl-piperazin-1-yl)-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((4-tert.butoxycarbonyl-piperazin-1-yl)-methylcarbonyl)-N-isopropyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{37}H_{43}N_5O_6$

ESI mass spectrum: $m/z = 652 [M-H^-]$

25

(117) 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-benzyl-p-phenylenediamine

30 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{40}H_{36}N_4O_4$

ESI mass spectrum: $m/z = 635$ $[M-H]^-$

(118) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 5 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and N-(dimethylaminomethyl-carbonyl)-N-benzyl-p-phenylenediamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{34}H_{32}N_4O_4$

ESI mass spectrum: $m/z = 559$ $[M-H]^-$

10

(119) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(5-methyl-imidazol-4-yl)-aniline

- 15 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{37}H_{36}N_4O_4$

ESI mass spectrum: $m/z = 559$ $[M-H]^-$

(120) 3-Z-[1-(4-(1,2,4-triazol-2-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(1,2,4-triazol-1-yl-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

$C_{26}H_{21}N_5O_3$

- 25 ESI mass spectrum: $m/z = 450$ $[M-H]^-$

(121) 3-Z-[1-(4-(1,2,3-triazol-2-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(1,2,3-triazol-2-yl-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 20:1)




ESI mass spectrum: $m/z = 450 [M-H^-]$

- 5 (122) 3-Z-[1-(4-(1,2,3-triazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(1,2,3-triazol-1-yl-methyl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/methanol = 9:1)



- 10 ESI mass spectrum: $m/z = 450 [M-H^-]$

- (123) 3-Z-[1-(4-((N-aminocarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((N-aminocarbonylmethyl-N-methyl-amino)-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)



ESI mass spectrum: $m/z = 469 [M-H^-]$

- 20 (124) 3-Z-[1-(4-((di-(2-methoxy-ethyl)-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((di-(2-methoxy-ethyl)-amino)-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

- 25 $C_{30}H_{33}N_3O_5$

ESI mass spectrum: $m/z = 514 [M-H^-]$

- (125) 3-Z-[1-(4-(pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 30 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(pyrrolidin-1-yl-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₂₈H₂₇N₃O₃

ESI mass spectrum: m/z = 452 [M-H⁻]

- 5 (126) 3-Z-[1-(4-((di-(2-hydroxy-ethyl)-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((di-(2-hydroxy-ethyl)-amino)-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

- 10 C₂₈H₂₉N₃O₅

ESI mass spectrum: m/z = 486 [M-H⁻]

(127) 3-Z-[1-(4-((N-ethoxycarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- 15 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((N-ethoxycarbonylmethyl-N-methyl-amino)-methyl)-aniline

R_f value: 0.5 (aluminium oxide, methylene chloride/ethanol = 40:1)

C₂₉H₂₉N₃O₅

ESI mass spectrum: m/z = 498 [M-H⁻]

- 20

(128) 3-Z-[1-(4-(azetidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(azetidin-1-yl-methyl)-aniline

- 25 R_f value: 0.5 (silica gel, methylene chloride/methanol/ammonia = 9:1:0.5)

C₂₇H₂₅N₃O₃

ESI mass spectrum: m/z = 438 [M-H⁻]

- 30 (129) 3-Z-[1-(4-(N-propyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(N-propyl-N-tert.butoxycarbonyl-aminomethyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₃₂H₃₅N₃O₅

5 ESI mass spectrum: m/z = 540 [M-H⁻]

(130) 3-Z-[1-(4((N-(2-(2-methoxy-ethoxy)-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
10 and 4-((N-(2-(2-methoxy-ethoxy)-ethyl)-N-methyl-amino)-methyl)-aniline

R_f value: 0.4 (silica gel, methylene chloride/methanol = 9:1)

C₃₀H₃₃N₃O₅

ESI mass spectrum: m/z = 514 [M-H⁻]

15 (131) 3-Z-[1-(4-((N-(tert.butoxycarbonyl-3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-(N-(N-tert.butoxycarbonyl-3-amino-propyl)-N-methyl-aminomethyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

20 C₃₃H₃₈N₄O₅

ESI mass spectrum: m/z = 571 [M+H⁺]

(132) 3-Z-[1-(4-((N-(methylcarbamoyl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone
and 4-((N-(methylcarbamoyl-methyl)-N-methyl-amino)-methyl)-aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

C₂₈H₂₈N₄O₄

ESI mass spectrum: m/z = 483 [M-H⁻]

30

(133) 3-Z-[1-(4-((N-(dimethylcarbamoyl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((N-(dimethylcarbamoyl-methyl)-N-methyl-amino)-methyl)-aniline

5 R_f value: 0.3 (silica gel, methylene chloride/methanol = 10:1)

$C_{29}H_{30}N_4O_4$

ESI mass spectrum: $m/z = 497 [M-H^-]$

(134) 3-Z-[1-(4-methyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-methyl-aniline

R_f value: 0.4 (silica gel, methylene chloride/methanol = 9:1)

$C_{24}H_{20}N_2O_3$

ESI mass spectrum: $m/z = 383 [M-H^-]$

15

(135) 3-Z-[1-(4-((N-propyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((N-propyl-N-methyl-amino)-methyl)-aniline

20 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{28}H_{29}N_3O_3$

ESI mass spectrum: $m/z = 454 [M-H^-]$

(136) 3-Z-[1-(4-((N-(2-hydroxy-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((N-(2-hydroxy-ethyl)-N-methyl-amino)-methyl)-aniline

R_f value: 0.5 (aluminium oxide, methylene chloride/ethanol = 40:1)

$C_{27}H_{27}N_3O_4$

30 ESI mass spectrum: $m/z = 456 [M-H^-]$

(137) 3-Z-[1-(4-((N-(2-dimethylamino-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((N-(2-dimethylamino-ethyl)-N-methyl-amino)-methyl)-aniline

5 R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{29}H_{32}N_4O_3$

ESI mass spectrum: $m/z = 483 [M-H^-]$

10 (138) 3-Z-[1-(4-((N-(3-dimethylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-((N-(3-dimethylamino-propyl)-N-methyl-amino)-methyl)-aniline aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

15 $C_{30}H_{34}N_4O_3$

ESI mass spectrum: $m/z = 497 [M-H^-]$

20 (139) 3-Z-[1-(4-(3-oxo-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 1-acetyl-3-(1-ethoxy-1-phenylmethylene)-6-methoxycarbonyl-2-indolinone and 4-(3-oxo-piperazin-1-yl-methyl)-aniline aniline

R_f value: 0.46 (silica gel, methylene chloride/methanol = 9:1)

$C_{28}H_{26}N_4O_4$

25 ESI mass spectrum: $m/z = 481 [M-H^-]$

Example 4

30 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone

485 mg of 3-Z-[1-(4-tert.butoxycarbonyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone are dissolved in 15 ml of methylene chloride and 6.0 ml of trifluoroacetic acid are added. The mixture is stirred for 2 hours at room temperature. Then the solvent is removed and the residue recrystallised from ether.

5 Yield: 375 mg (87 % of theory),

R_f value: 0.3 (silica gel, methylene chloride/methanol = 10:1)

C₂₅H₂₀N₂O₅

Mass spectrum: m/z = 428 [M⁺]

10 The following compounds are prepared analogously to Example 4:

(1) 3-Z-[1-(4-aminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
Prepared from 3-Z-[1-(4-(N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 R_f value: 0.5 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)

C₂₄H₂₁N₃O₃

ESI mass spectrum: m/z = 398 [M-H⁻]

(2) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-tert.butoxycarbonyl-ethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.4 (silica gel, methylene chloride/methanol/ammonia = 10:1:0.01)


25 C₂₆H₂₅N₃O₃

ESI mass spectrum: m/z = 426 [M-H⁻]

(3) 3-Z-[1-(4-carboxymethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
Prepared from 3-Z-[1-(4-tert.butoxycarbonylmethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30 ethoxycarbonyl-2-indolinone

R_f value: 0.1 (aluminium oxide, methylene chloride/ethanol/ammonia = 5:1:0.01)



ESI mass spectrum: $m/z = 441$ $[M-H]^-$

(4) 3-Z-[1-(4-carboxy-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 Prepared from 3-Z-[1-(4-tert.butoxycarbonyl-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

R_f value: 0.1 (aluminium oxide, methylene chloride/ethanol = 20:1)



ESI mass spectrum: $m/z = 379$ $[M-H]^-$

10

(5) 3-Z-[1-(4-(piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(4-tert.butoxycarbonyl-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 R_f value: 0.1 (silica gel, methylene chloride/methanol/ammonia = 10:1:0.01)




ESI mass spectrum: $m/z = 469$ $[M+H]^+$

(6) 3-Z-[1-(4-butylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20

Prepared from 3-Z-[1-(4-(N-butyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)




25 ESI mass spectrum: $m/z = 454$ $[M-H]^-$

(7) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

Prepared from 3-Z-[1-(4-(N-tert.butoxycarbonyl-N-ethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

R_f value: 0.3 (silica gel, methylene chloride/methanol/ammonia = 10:1:0.01)




ESI mass spectrum: $m/z = 442 [M+H^+]$

(8) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

5 Prepared from 3-Z-[1-(4-(N-tert.butoxycarbonyl-ethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

R_f value: 0.2 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)



ESI mass spectrum: $m/z = 411 [M-H^-]$


10

(9) 3-Z-[1-(4-(N-(piperazin-1-yl-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-((4-tert.butoxycarbonyl-piperazin-1-yl)-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15

R_f value: 0.35 (silica gel, methylene chloride/methanol = 9:1)




ESI mass spectrum: $m/z = 552 [M-H^-]$

20

(10) 3-Z-[1-(4-(N-((2-(piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-((2-(4-tert.butoxycarbonyl-piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 R_f value: 0.4 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)



ESI mass spectrum: $m/z = 540 [M+H^+]$

(11) 3-Z-[1-(4-(N-propyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-

30 methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-propyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.35 (silica gel, methylene chloride/methanol = 9:1)

C₂₇H₂₇N₃O₃

5 ESI mass spectrum: m/z = 440 [M-H⁻]

(12) 3-Z-[1-(4-((N-(3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-((N-(tert.butoxycarbonyl-3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10

R_f value: 0.35 (silica gel, methylene chloride/methanol = 9:1)

C₂₈H₃₀N₄O₃

ESI mass spectrum: m/z = 471 [M+H⁺]

15 Example 5

3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 100 mg of 3-Z-[1-(4-(N-benzyl-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone are dissolved in 20 ml of ethanol, 0.2 ml of 1N hydrochloric acid are added and the mixture is hydrogenated for 70 minutes at room temperature and 50 psi hydrogen pressure. The reaction solution is filtered and the filtrate concentrated by rotary evaporation. The residue is dried *in vacuo* at 100°C.

25 Yield: 50 mg (53 % of theory),

R_f value: 0.3 (silica gel, methylene chloride/ethanol/ammonia = 5:1:0.01)

C₂₆H₂₅N₃O₃

ESI mass spectrum: m/z = 426 [M-H⁻]

30 The following compounds are prepared analogously to Example 5:

(1) 3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-benzyl-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 R_f value: 0.2 (silica gel, methylene chloride/methanol/ammonia = 10:1:0.01)

$C_{25}H_{23}N_3O_3$

ESI mass spectrum: $m/z = 412$ [M-H]

(2) 3-Z-[1-(4-(N-(2-methylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 Prepared from 3-Z-[1-(4-((2-(N-benzyl-N-methyl-amino)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.3 (silica gel, methylene chloride/methanol/ammonia = 10:1:0.01)

$C_{27}H_{28}N_4O_5S$

15 ESI mass spectrum: $m/z = 519$ [M-H]

(3) 3-Z-[1-(4-(N-(2-amino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 Prepared from 3-Z-[1-(4-(N-cyanomethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.5 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)

$C_{26}H_{26}N_4O_5S$

ESI mass spectrum: $m/z = 505$ [M-H]

25 (4) 3-Z-[1-(4-(N-(3-methylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-(3-(N-benzyl-N-methyl-amino)-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.3 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)

30 $C_{28}H_{30}N_4O_5S$

ESI mass spectrum: $m/z = 533$ [M-H]

(5) 3-Z-[1-(4-(N-(piperazin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-((4-benzyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.5 (silica gel, methylene chloride/methanol = 9:1)

$C_{30}H_{31}N_5O_4$

ESI mass spectrum: $m/z = 524 [M-H^-]$

(6) 3-Z-[1-(4-(N-(methylamino-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.3 (silica gel, methylene chloride/methanol = 9:1)

$C_{27}H_{26}N_4O_4$

ESI mass spectrum: $m/z = 469 [M-H^-]$

(7) 3-Z-[1-(4-(N-((2-methylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-((2-(N-benzyl-N-methyl-amino)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.3 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)

$C_{28}H_{28}N_4O_4$

ESI mass spectrum: $m/z = 483 [M-H^-]$

Example 6

3-Z-[1-(4-ureidomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

300 mg of 3-Z-[1-(4-aminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone are dissolved in 15 ml of methanol and 200 ml of triethylamine are added. Then

400 mg of potassium cyanate in 5 ml of water are added. After 2 days of stirring at room temperature the reaction solution is concentrated by rotary evaporation, the residue taken up in methylene chloride and washed once with water and once with saturated sodium chloride solution. The organic phase is dried over sodium sulphate and concentrated by rotary evaporation. The residue is dried *in vacuo* at 100°C.

Yield: 100 mg of (21 % of theory),

R_f value: 0.7 (silica gel, methylene chloride/methanol = 5:1)

C₂₅H₂₂N₄O₄

ESI mass spectrum: m/z = 441 [M-H]

Example 7

3-Z-[1-(4-guanidinomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

300 mg of 3-Z-[1-(4-aminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone are dissolved in 5 ml of dimethylformamide and 300 ml of triethylamine are added. Then 700 mg of 3,5-dimethylpyrazol-1-carboxylic acid amidine in 5 ml of dimethylformamide are added. After one day of stirring at room temperature the reaction solution is concentrated by rotary evaporation. The residue is dried at 100°C *in vacuo*.

Yield: 200 mg (87 % of theory),

R_f value: 0.1 (Reversed phase RP 8, methanol/five percent saline solution = 6:4)

C₂₅H₂₃N₅O₃

Mass spectrum: m/z = 441 [M⁺]

Example 8

3-Z-[1-(4-acetylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

100 mg of 3-Z-[1-(4-aminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone are dissolved in 5 ml of glacial acetic acid, 0.1 ml of acetic anhydride is added and the mixture is stirred for 10 minutes at room temperature. After this time the reaction solution is poured onto saturated soda solution and extracted four times with methylene chloride. The combined organic phases are washed with saturated saline solution, dried over sodium sulphate and concentrated by rotary evaporation. The residue is dried at 100°C *in vacuo*.

Yield: 20 mg (23 % of theory),

R_f value: 0.4 (silica gel, methylene chloride/methanol = 10:1)

10 C₂₆H₂₃N₃O₄

ESI mass spectrum: m/z = 440 [M-H⁺]

The following compounds are prepared analogously to Example 8:

15 (1) 3-Z-[1-(4-(N-methylsulphonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-aminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone and methanesulphonyl chloride/triethylamine

R_f value: 0.7 (silica gel, methylene chloride/methanol = 5:1)

20 C₂₅H₂₃N₃O₅S

ESI mass spectrum: m/z = 476 [M-H⁺]

(2) 3-Z-[1-(4-(4-benzoyl-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 Prepared from 3-Z-[1-(4-(piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone and benzoyl chloride

R_f value: 0.7 (silica gel, methylene chloride/methanol = 10:1)

C₃₅H₃₂N₄O₄

ESI mass spectrum: m/z = 571 [M-H⁺]

30

(3) 3-Z-[1-(4-((N-(3-acetylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-((N-(3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 R_f value: 0.3 (silica gel, methylene chloride/methanol = 9:1)

$C_{30}H_{32}N_4O_4$

ESI mass spectrum: $m/z = 511 [M-H^-]$

Example 9

10

3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

0.8 g of 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone are dissolved in 30 ml of ethanol, 8.3 ml of 1N sodium hydroxide solution are added and the mixture is stirred for 1 hour at 80°C. After cooling, it is neutralised with 8.3 ml of 1N hydrochloric acid. The precipitate formed is suction filtered, washed with water, ethanol and ether and dried *in vacuo* at 100°C.

15 Yield: 0.7 g of (89 % of theory),

R_f value: 0.2 (silica gel, methylene chloride/methanol = 5:2)

20 $C_{28}H_{27}N_3O_3$

Mass spectrum: $m/z = 453 [M^+]$

The following compounds are prepared analogously to Example 9:

25 (1) 3-Z-[1-(4-bromo-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

Prepared from 3-Z-[1-(4-bromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

R_f value: 0.4 (silica gel, toluene/ethyl acetate = 5:1)

$C_{22}H_{15}BrN_2O_3$

30 ESI mass spectrum: $m/z = 435/437 [M+H^+]$

(2) 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

Prepared from 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 R_f value: 0.7 (Reversed phase RP 8, methanol/five percent saline solution = 4:1)

$C_{25}H_{23}N_3O_3$

ESI mass spectrum: $m/z = 414 [M+H^+]$

10 (3) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

Prepared from 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

R_f value: 0.7 (Reversed phase RP 8, methanol/five percent saline solution = 4:1)

$C_{25}H_{23}N_3O_3$

15 ESI mass spectrum: $m/z = 412 [M-H^-]$

(4) 3-Z-[1-(4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

20 Prepared from 3-Z-[1-(4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

R_f value: 0.6 (Reversed phase RP 8, methanol/five percent saline solution = 4:1)

$C_{30}H_{31}N_3O_3$

ESI mass spectrum: $m/z = 482 [M+H^+]$

25 (5) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

Prepared from 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.6 (Reversed phase RP 8, methanol/five percent saline solution = 4:1)

30 $C_{26}H_{20}N_4O_3$

ESI mass spectrum: $m/z = 435 [M-H^-]$

(6) 3-Z-[1-(4-(N-acetyl-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

Prepared from 3-Z-[1-(4-(N-acetyl-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.3 (silica gel, methylene chloride/methanol = 10:1)

C₂₈H₂₆N₄O₅

ESI mass spectrum: m/z = 497 [M-H⁻]

10 (7) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

Prepared from 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.6 (Reversed phase RP 8, methanol/five percent saline solution = 4:1)

C₂₅H₂₃N₃O₃

15 ESI mass spectrum: m/z = 412 [M-H⁻]

(8) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

Prepared from 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

R_f value: 0.6 (Reversed phase RP 8, methanol/five percent saline solution = 4:1)

C₂₇H₂₆N₄O₄

ESI mass spectrum: m/z = 469 [M-H⁻]

25 (9) 3-Z-[1-(4-(N-tert.butoxycarbonyl-ethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

Prepared from 3-Z-[1-(4-(N-tert.butoxycarbonyl-ethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

R_f value: 0.4 (silica gel, methylene chloride/methanol = 10:1)

30 C₃₀H₃₁N₃O₅

ESI mass spectrum: m/z = 512 [M-H⁻]

(10) 3-Z-[1-(4-((N-carboxymethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-((N-ethoxycarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl -2-indolinone

R_f value: 0.4 (silica gel, methylene chloride/methanol = 6:1)

C₂₇H₂₅N₃O₅

ESI mass spectrum: m/z = 470 [M-H⁻]

Example 10

3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

0.9 g of 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone are suspended in 35 ml of dimethylformamide and 0.4 g of carbonyldiimidazole are added. The mixture is stirred for 14 hours at 80°C. After this time 20 ml of methanol are added and the mixture is stirred for another 3 hours at 50°C. The solvent is removed and the residue is purified over a silica gel column with methylene chloride/methanol (3:1) as eluant.

Yield: 0.5 g of (49% of theory),

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 30:1)

C₂₉H₂₉N₃O₃

ESI mass spectrum: m/z = 468 [M+H⁺]

The following compounds are prepared analogously to Example 10:

(1) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-benzyloxycarbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and benzyl alcohol

R_f value: 0.6 (aluminium oxide, methylene chloride/methanol = 30:1)

C₃₅H₃₃N₃O₃

5 Mass spectrum: m/z = 543 [M⁺]

(2) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-isopropoxy carbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and isopropanol

R_f value: 0.4 (aluminium oxide, methylene chloride/isopropanol = 30:1)

C₃₁H₃₃N₃O₃

Mass spectrum: m/z = 495 [M⁺]

15

(3) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-propyloxy carbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and n-propanol

20 R_f value: 0.7 (silica gel, methylene chloride/methanol = 5:1)

C₃₁H₃₃N₃O₃

Mass spectrum: m/z = 495 [M⁺]

25 (4) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-butyloxy carbonyl-2-indolinone

Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and n-butanol

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

C₃₂H₃₅N₃O₃

30 Mass spectrum: m/z = 509 [M⁺]

(5) 3-Z-[1-(4-bromo-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from 3-Z-[1-(4-bromo-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and ammonia

R_f value: 0.5 (silica gel, methylene chloride/methanol = 10:1)

5 C₂₂H₁₆BrN₂O₃

Mass spectrum: m/z = 432/434 [M-H⁻]

(6) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

10 Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and ethylamine gas

R_f value: 0.6 (silica gel, methylene chloride/methanol = 5:1)

C₃₀H₃₂N₄O₂

Mass spectrum: m/z = 480 [M⁺]

15

(7) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(2-methoxy-ethoxy)-carbonyl]-2-indolinone

Prepared from 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and methylglycol

20 R_f value: 0.8 (silica gel, methylene chloride/methanol = 4:1)

C₂₅H₂₃N₃O₃

ESI mass spectrum: m/z = 470 [M-H⁻]

(8) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(2-dimethylamino-ethoxy)-carbonyl]-2-indolinone

25

Prepared from 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and 2-dimethylaminoethanol aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 5:2)

C₂₉H₃₂N₄O₃

30 ESI mass spectrum: m/z = 483 [M-H⁻]

(9) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(2-N-tert.butoxycarbonyl-amino-ethoxy)-carbonyl]-2-indolinone

Prepared from 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and 2-N-tert.butoxycarbonyl-amino-ethanol aniline

5 R_f value: 0.8 (silica gel, methylene chloride/methanol = 5:2)

$C_{32}H_{36}N_4O_5$

ESI mass spectrum: $m/z = 412 [M-H^-]$

10 (10) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(2,2,2-trifluoroethoxy)-carbonyl]-2-indolinone

Prepared from 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and 2,2,2-trifluoroethanol aniline

R_f value: 0.5 (silica gel, methylene chloride/methanol = 5:1)

$C_{27}H_{24}F_3N_3O_3$

15 ESI mass spectrum: $m/z = 494 [M-H^-]$

Example 11

3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

20

0.9 g of 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone, 0.8 g of TBTU and 0.4 g of HOBT are suspended in 25 ml of dimethylformamide and 1.0 ml of triethylamine are added. The mixture is stirred for 15 minutes at room temperature. After this time ammonia gas is introduced at 10-15°C over a period of 15 minutes and the mixture is stirred for 1.5 hours at room temperature. The precipitate formed is suction filtered, washed with water, ethanol and ether and dried at 100°C *in vacuo*.

25

Yield: 0.6 g (64 % of theory),

R_f value: 0.4 (Reversed phase RP 8, methanol/five percent saline solution = 6:4)

30

$C_{28}H_{28}N_4O_2$

ESI mass spectrum: $m/z = 453 [M+H^+]$

The following compounds are prepared analogously to Example 11:

- (1) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-dimethylcarbamoyl-
5 2-indolinone

Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-
2-indolinone and dimethylamine hydrochloride/diisopropylethylamine

R_f value: 0.5 (silica gel, methylene chloride/methanol = 5:1)

C₃₀H₃₂N₄O₂

- 10 ESI mass spectrum: m/z = 481 [M+H⁺]

- (2) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-(N-ethyl-N-methyl-
carbamoyl)-2-indolinone

- Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-
15 2-indolinone and N-ethyl-N-methyl-amine

R_f value: 0.5 (aluminium oxide, methylene chloride/ethanol = 20:1)

C₃₁H₃₄N₄O₂

ESI mass spectrum: m/z = 495 [M+H⁺]

- 20 (3) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methylcarbamoyl-2-
indolinone

Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-
2-indolinone and methylamine hydrochloride/diisopropylethylamine

R_f value: 0.3 (aluminium oxide, methylene chloride/ethanol = 20:1)

- 25 C₂₉H₃₀N₄O₂

ESI mass spectrum: m/z = 467 [M+H⁺]

- (4) 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methylcarbamoyl-2-
indolinone

- 30 Prepared from 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-
carboxyl-2-indolinone and methylamine hydrochloride/triethylamine

R_f value: 0.3 (silica gel, methylene chloride/ethanol = 2:1)

C₂₆H₂₆N₄O₂

Mass spectrum: m/z = 426 [M⁺]

- 5 (5) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-(2-hydroxyethyl-carbamoyl)-2-indolinone

Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and ethanolamine/diisopropylethylamine

R_f value: 0.5 (aluminium oxide, methylene chloride/methanol = 20:1)

- 10 C₃₀H₃₂N₄O₃

ESI mass spectrum: m/z = 495 [M-H⁻]

(6) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-diethylcarbamoyl-2-indolinone

- 15 Prepared from 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone and diethylamine hydrochloride/diisopropylethylamine

R_f value: 0.8 (aluminium oxide, methylene chloride/methanol = 10:1)

C₃₂H₃₆N₄O₂

ESI mass spectrum: m/z = 509 [M+H⁺]

- 20

(7) 3-Z-[1-(4-(N-tert.butoxycarbonyl-ethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-tert.butoxycarbonyl-ethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

- 25 R_f value: 0.3 (silica gel, toluene/ethyl acetate/ethanol = 4:2:1)

C₃₀H₃₂N₄O₄

ESI mass spectrum: m/z = 511 [M-H⁻]

- 30 (8) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

Prepared from 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

R_f value: 0.5 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)

5 C₂₇H₂₇N₅O₃

ESI mass spectrum: m/z = 468 [M-H⁻]

Example 12

10 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone x citric acid

3.25 g of citric acid monohydrate are placed in 50 ml of methanol and 5.0 g of 3-Z-[1-(4-(N-dimethylaminomethyl-carbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone are added at room temperature. The solution formed is evaporated down, the residue is washed with ether and recrystallised from ethyl acetate. Yield: 6.3 g (90 % of theory),

R_f value: 0.6 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)

Melting point: 198°C

20 C₂₈H₂₈N₄O₅ x C₆H₈O₇

ESI mass spectrum: m/z = 483 [M-H⁻]

Elemental analysis: calc.: C 60.34 H 5.37 N 8.28

found: 59.98 5.25 8.13

25 The following compound is prepared analogously to Example 12:

(1) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone x methanesulphonic acid

Prepared from 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-

30 methoxycarbonyl-2-indolinone and methanesulphonic acid

R_f value: 0.6 (silica gel, methylene chloride/methanol/ammonia = 5:1:0.01)

Melting point: 275°C

$C_{26}H_{25}N_3O_3 \times CH_4O_3S$

ESI mass spectrum: $m/z = 426 [M-H^-]$

Elemental analysis: calc.: C 61.92 H 5.59 N 8.03 S 6.12

5 found: 61.43 5.87 7.85 5.39

The following compounds may be prepared analogously to the foregoing Examples:

- 10 (1) 3-Z-(1-anilino-1-phenyl-methylene)-6-ethoxycarbonyl-2-indolinone
- (2) 3-Z-[1-(4-nitro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (3) 3-Z-[1-(4-fluoro-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone
- 15 (4) 3-Z-[1-(4-chloro-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone
- (5) 3-Z-[1-(4-iodo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (6) 3-Z-[1-(4-cyano-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone
- 20 (7) 3-Z-[1-(4-methoxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (8) 3-Z-[1-(4-ethoxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 25 (9) 3-Z-[1-(4-trifluoromethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (10) 3-Z-[1-(4-methyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 30 (11) 3-Z-[1-(4-methylmercapto-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(12) 3-Z-[1-(4-aminomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(13) 3-Z-[1-(4-(isopropylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(14) 3-Z-[1-(4-(anilinomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(15) 3-Z-[1-(4-(propylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(16) 3-Z-[1-(4-(butylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(17) 3-Z-[1-(4-(isobutylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(18) 3-Z-[1-(4-(cyclohexylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(19) 3-Z-[1-(4-(benzylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(20) 3-Z-[1-(4-((N-ethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(21) 3-Z-[1-(4-((N-methyl-N-propyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(22) 3-Z-[1-(4-((N-isopropyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (23) 3-Z-[1-(4-((N-ethyl-N-propyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(24) 3-Z-[1-(4-((N-ethyl-N-isopropyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (25) 3-Z-[1-(4-(dipropylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(26) 3-Z-[1-(4-(diisopropylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (27) 3-Z-[1-(4-((N-benzyl-N-ethyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (28) 3-Z-[1-(4-(dibenzylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(29) 3-Z-[1-(4-(3,6-dihydro-2H-pyridin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (30) 3-Z-[1-(4-(3,5-dimethyl-piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(31) 3-Z-[1-(4-(azepan-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(32) 3-Z-[1-(4-(piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (33) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(34) 3-Z-[1-(4-(thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (35) 3-Z-[1-(4-(1-oxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(36) 3-Z-[1-(4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (37) 3-Z-[1-(4-(acetylamino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (38) 3-Z-[1-(4-(2-amino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(39) 3-Z-[1-(4-(2-methylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (40) 3-Z-[1-(4-(2-ethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(41) 3-Z-[1-(4-(2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(42) 3-Z-[1-(4-(2-piperidin-1-yl-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (43) 3-Z-[1-(4-(2-acetylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(44) 3-Z-[1-(4-(3-amino-propyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (45) 3-Z-[1-(4-(3-dimethylamino-propyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(46) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (47) 3-Z-[1-(4-(N-methylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (48) 3-Z-[1-(4-(N-ethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(49) 3-Z-[1-(4-(N-diethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (50) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(51) 3-Z-[1-(4-(N-(morpholin-4-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(52) 3-Z-[1-(4-(N-(piperazin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (53) 3-Z-[1-(4-(N-(2-amino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(54) 3-Z-[1-(4-(N-(2-methylamino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (55) 3-Z-[1-(4-(N-(2-diethylamino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(56) 3-Z-[1-(4-(N-acetyl-N-(2-aminoethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (57) 3-Z-[1-(4-(N-acetyl-N-(2-methylamino-ethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (58) 3-Z-[1-(4-(N-acetyl-N-(2-methylamino-propyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(59) 3-Z-[1-(4-(N-acetyl-N-(2-piperidin-1-yl-ethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (60) 3-Z-[1-(4-(N-acetyl-N-(aminocarbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(61) 3-Z-[1-(4-(N-acetyl-N-(dimethylaminocarbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(62) 3-Z-[1-(4-(N-acetyl-N-(piperidin-1-yl-carbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (63) 3-Z-[1-(4-(N-methyl-N-(aminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(64) 3-Z-[1-(4-(N-methyl-N-(methylaminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (65) 3-Z-[1-(4-(N-methyl-N-(dimethylaminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(66) 3-Z-[1-(4-(N-methyl-N-(piperidin-1-yl-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (67) 3-Z-[1-(4-(N-(2-aminoethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (68) 3-Z-[1-(4-(N-(2-methylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(69) 3-Z-[1-(4-(N-(2-ethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (70) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30 (71) 3-Z-[1-(4-(N-(2-pyrrolidin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(72) 3-Z-[1-(4-(N-(2-piperidin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (73) 3-Z-[1-(4-(N-(2-piperazin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(74) 3-Z-[1-(4-(N-(2-(morpholin-4-yl)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (75) 3-Z-[1-(4-(N-(aminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(76) 3-Z-[1-(4-(N-(methylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (77) 3-Z-[1-(4-(N-(ethylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (78) 3-Z-[1-(4-(N-(N-(2-dimethylamino-ethyl)-N-methyl-amino)-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(79) 3-Z-[1-(4-(N-(diethylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (80) 3-Z-[1-(4-(N-(pyrrolidin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(81) 3-Z-[1-(4-(N-(piperidin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(82) 3-Z-[1-(4-(N-(piperazin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(83) 3-Z-[1-(4-(N-((morpholin-4-yl)-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(84) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(85) 3-Z-[1-(4-(3-dimethylamino-propoxy)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(86) 3-Z-[1-(4-(aminocarbonylmethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(87) 3-Z-[1-(4-(2-aminocarbonyl-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(88) 3-Z-[1-(4-(pyridin-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(89) 3-Z-[1-(4-(pyridine-3-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(90) 3-Z-[1-(4-(pyridin-4-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(91) 3-Z-[1-(4-(N-acetyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(92) 3-Z-[1-(4-(N-ethylcarbonyl-N-(dimethylaminocarbonyl-methyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(93) 3-Z-[1-(carbamoylmethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (94) 3-Z-[1-(4-dimethylcarbamoylmethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(95) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

10 (96) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(97) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

15 (98) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(99) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-ethylidene]-6-ethoxycarbonyl-2-indolinone

20 (100) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(101) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

25 (102) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

30 (103) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

- (104) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone
- (105) 3-Z-[1-(4-tetrazol-5-yl-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone
- 5 (106) 3-Z-[1-(4-tetrazol-5-yl-anilino)-ethylidene]-6-ethoxycarbonyl-2-indolinone
- (107) 3-Z-[1-(4-tetrazol-5-yl-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone
- 10 (108) 3-Z-[1-(4-tetrazol-5-yl-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone
- (109) 3-Z-[1-(4-carboxy-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone
- (110) 3-Z-[1-(4-carboxy-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone
- 15 (111) 3-Z-[1-(4-carboxy-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone
- (112) 3-Z-[1-(4-(N-(3-dimethylamino-propionyl)-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 20 (113) 3-Z-[1-(4-(N-(4-dimethylamino-butyryl)-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (114) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-(2-dimethylamino-ethylsulphonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 25 (115) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-(3-dimethylamino-propylsulphonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 30 (116) 3-Z-[1-(4-((2-hydroxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

- (117) 3-Z-[1-(4-((2-methoxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 5 (118) 3-Z-[1-(4-((2-dimethylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (119) 3-Z-[1-(4-((3-dimethylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 10 (120) 3-Z-[1-(4-((N-tert.butoxycarbonyl-2-amino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (121) 3-Z-[1-(4-((N-tert.butoxycarbonyl-3-amino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 15 (122) 3-Z-[1-(4-((2-amino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (123) 3-Z-[1-(4-((3-amino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 20 (124) 3-Z-[1-(4-((2-acetylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 25 (125) 3-Z-[1-(4-((3-acetylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- (126) 3-Z-[1-(4-((2-methylsulphonylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone
- 30

(127) 3-Z-[1-(4-((3-methylsulphonylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (128) 3-Z-[1-(4-(N-(N-tert.butoxycarbonyl-2-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(129) 3-Z-[1-(4-(N-(2-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (130) 3-Z-[1-(4-(N-(2-acetylamino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(131) 3-Z-[1-(4-(N-(2-methylsulphonylamino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (132) 3-Z-[1-(4-(carboxymethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (133) 3-Z-[1-(4-(ethoxycarbonylmethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(134) 3-Z-[1-(4-(carbamoylemethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (135) 3-Z-[1-(4-(dimethylcarbamoylemethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(136) 3-Z-[1-(4-(methylcarbamoylemethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(137) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (138) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-nitro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(139) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-acetylamino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (140) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methylsulphonylamino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(141) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-cyano-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (142) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-hydroxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (143) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methoxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(144) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-ethoxycarbonyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (145) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-carboxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(146) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-carbamoyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(147) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-chloro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (148) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-fluoro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(149) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-bromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (150) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(151) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-trifluoromethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (152) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3,5-dibromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (153) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3,5-dichloro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(154) 3-Z-[1-(4-(dimethylaminomethyl)-3-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (155) 3-Z-[1-(4-(dimethylaminomethyl)-3-nitro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(156) 3-Z-[1-(4-(dimethylaminomethyl)-3-acetyl-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(157) 3-Z-[1-(4-(dimethylaminomethyl)-3-(methylsulphonylamino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (158) 3-Z-[1-(4-(dimethylaminomethyl)-3-cyano-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(159) 3-Z-[1-(4-(dimethylaminomethyl)-3-hydroxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (160) 3-Z-[1-(4-(dimethylaminomethyl)-3-methoxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(161) 3-Z-[1-(4-(dimethylaminomethyl)-3-(ethoxycarbonyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (162) 3-Z-[1-(4-(dimethylaminomethyl)-3-carboxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (163) 3-Z-[1-(4-(dimethylaminomethyl)-3-carbamoyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(164) 3-Z-[1-(4-(dimethylaminomethyl)-3-chloro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (165) 3-Z-[1-(4-(dimethylaminomethyl)-3-fluoro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(166) 3-Z-[1-(4-(dimethylaminomethyl)-3-bromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(167) 3-Z-[1-(4-(dimethylaminomethyl)-3-methyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(168) 3-Z-[1-(4-(dimethylaminomethyl)-3-trifluoromethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(169) 3-Z-[1-(4-(dimethylaminomethyl)-3,5-dibromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(170) 3-Z-[1-(4-(dimethylaminomethyl)-3,5-dichloro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(171) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(172) 3-Z-[1-(4-(N-(imidazo-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(173) 3-Z-[1-(4-(N-(phthalimido-2-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(174) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(175) 3-Z-[1-(4-(N-acetylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(176) 3-Z-[1-(4-(N-methylsulphonylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(177) 3-Z-[1-(4-(N-((N-(2-methoxyethyl)-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(178) 3-Z-[1-(4-(N-((N-(2-dimethylaminoethyl)-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(179) 3-Z-[1-(4-(N-((di-(2-hydroxyethyl)-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(180) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(181) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-ethylidene]-6-ethoxycarbonyl-2-indolinone

(182) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(183) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(184) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(185) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-ethylidene]-6-ethoxycarbonyl-2-indolinone

(186) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(187) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(188) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-
5 6-ethoxycarbonyl-2-indolinone

(189) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (190) 3-Z-[1-(4-((imidazolidin-2,4-dion-5-ylidene)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(191) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (192) 3-Z-[1-(4-(N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(193) 3-Z-[1-(4-(2-oxo-pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-
20 ethoxycarbonyl-2-indolinone

(194) 3-Z-[1-(4-(N-aminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (195) 3-Z-[1-(4-(N-cyanomethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(196) 3-Z-[1-(4-(2-(imidazol-4-yl)-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(197) 3-Z-[1-(4-((2-(N-benzyl-N-methyl-amino)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (198) 3-Z-[1-(4-cyclohexylamino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(199) 3-Z-[1-(4-(imidazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (200) 3-Z-[1-(4-(imidazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(201) 3-Z-[1-(N-methyl-piperidine-4-yl-amino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (202) 3-Z-[1-(4-(imidazol-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (203) 3-Z-[1-(4-((4-hydroxy-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(204) 3-Z-[1-(4-((4-methoxy-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (205) 3-Z-[1-(4-benzyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(206) 3-Z-[1-(4-(N-(3-trifluoroacetyl-amino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30 (207) 3-Z-[1-(4-(4-tert.butoxycarbonyl-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(208) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (209) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(210) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10

(211) 3-Z-[1-(4-((3-(N-benzyl-N-methyl-amino)-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (212) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(213) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-butyryl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (214) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-isobutyryl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(215) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-benzoyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25

(216) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30 (217) 3-Z-[1-(4-(4-hydroxymethyl-piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(218) 3-Z-[1-(4-(2-(4-hydroxy-piperidin-1-yl)-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (219) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-propylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(220) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-butylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (221) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-phenylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(222) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-benzylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (223) 3-Z-[1-(4-((imidazolidin-2,4-dion-5-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (224) 3-Z-[1-(4-((3-hydroxy-pyrrolidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(225) 3-Z-[1-(4-(cyclohexyl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (226) 3-Z-[1-(4-(cyclohexyl-carbonyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(227) 3-Z-[1-(4-diethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(228) 3-Z-[1-(4-(N-(n-hexyl)-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (229) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(furan-2-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(230) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(2-methoxy-benzoyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (231) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(pyridine-3-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(232) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(phenyl-acetyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (233) 3-Z-[1-(4-(imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (234) 3-Z-[1-(4-(1-ethyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(235) 3-Z-[1-(4-(1-benzyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (236) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-isopropylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30 (237) 3-Z-[1-(4-(N-((4-benzyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(238) 3-Z-[1-(4-(N-(pyrrolidin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(239) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-bromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(240) 3-Z-[1-(4-(5-methyl-imidazol-4-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(241) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(242) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(243) 3-Z-[1-(4-(N-butyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(244) 3-Z-[1-(4-(N-((N-aminocarbonylmethyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(245) 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(246) 3-Z-[1-(4-(N-(di-(2-methoxyethyl)-amino-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(247) 3-Z-[1-(4-(N-((2-(4-tert.butoxycarbonyl-piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(248) 3-Z-[1-(4-(N-((2-(piperidin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (249) 3-Z-[1-(4-(N-((2-(N-benzyl-N-methyl-amino)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(250) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (251) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(252) 3-Z-[1-(4-(N-((4-tert.butoxycarbonyl-piperazin-1-yl)-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (253) 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (254) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(255) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (256) 3-Z-[1-(4-(1,2,4-triazol-2-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(257) 3-Z-[1-(4-(1,2,3-triazol-2-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(258) 3-Z-[1-(4-(1,2,3-triazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(259) 3-Z-[1-(4-((N-aminocarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(260) 3-Z-[1-(4-((di-(2-methoxy-ethyl)-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(261) 3-Z-[1-(4-((di-(2-hydroxy-ethyl)-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(262) 3-Z-[1-(4-((N-ethoxycarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(263) 3-Z-[1-(4-(azetidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(264) 3-Z-[1-(4-(N-propyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(265) 3-Z-[1-(4-((N-(2-(2-methoxy-ethoxy)-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(266) 3-Z-[1-(4-((N-(tert.butoxycarbonyl-3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(267) 3-Z-[1-(4-((N-(methylcarbamoyl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(268) 3-Z-[1-(4-((N-(dimethylcarbamoyl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

5 (269) 3-Z-[1-(4-((N-propyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(270) 3-Z-[1-(4-((N-(2-dimethylamino-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

10 (271) 3-Z-[1-(4-((N-(3-dimethylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(272) 3-Z-[1-(4-((N-(2-methoxy-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

15 (273) 3-Z-[1-(4-((N-(2-hydroxy-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

20 (274) 3-Z-[1-(4-((N-(dioxolan-2-yl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(275) 3-Z-[1-(4-(3-oxo-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

25 (276) 3-Z-[1-(4-(N-(piperazin-1-yl-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(277) 3-Z-[1-(4-(N-((2-(piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

30

(278) 3-Z-[1-(4-((N-(3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(279) 3-Z-[1-(4-(N-(3-methylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(280) 3-Z-[1-(4-Ureidomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(281) 3-Z-[1-(4-guanidinomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(282) 3-Z-[1-(4-(N-methylsulphonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(283) 3-Z-[1-(4-(4-benzoyl-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(284) 3-Z-[1-(4-((N-(3-acetylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(285) 3-Z-[1-(4-((N-(3-methylsulphonylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(286) 3-Z-[1-(4-((N-carboxymethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(287) 3-Z-(1-anilino-1-phenyl-methylene)-6-methoxycarbonyl-2-indolinone

(288) 3-Z-[1-(4-nitro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- (289) 3-Z-[1-(4-fluoro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (290) 3-Z-[1-(4-chloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 5 (291) 3-Z-[1-(4-bromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (292) 3-Z-[1-(4-iodo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (293) 3-Z-[1-(4-cyano-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 10 (294) 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (295) 3-Z-[1-(4-methoxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 15 (296) 3-Z-[1-(4-ethoxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (297) 3-Z-[1-(4-trifluoromethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 20 (298) 3-Z-[1-(4-methylmercapto-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (299) 3-Z-[1-(4-(isopropylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 25 (300) 3-Z-[1-(4-(anilinomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (301) 3-Z-[1-(4-(isobutylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 30

(302) 3-Z-[1-(4-(cyclohexylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (303) 3-Z-[1-(4-(benzylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(304) 3-Z-[1-(4-((N-methyl-N-propyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (305) 3-Z-[1-(4-((N-isopropyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(306) 3-Z-[1-(4-((N-ethyl-N-propyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (307) 3-Z-[1-(4-((N-ethyl-N-isopropyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (308) 3-Z-[1-(4-(dipropylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(309) 3-Z-[1-(4-(diisopropylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (310) 3-Z-[1-(4-((N-benzyl-N-ethyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(311) 3-Z-[1-(4-(dibenzylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(312) 3-Z-[1-(4-(3,6-dihydro-2H-pyridin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (313) 3-Z-[1-(4-(3,5-dimethyl-piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(314) 3-Z-[1-(4-(azepan-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (315) 3-Z-[1-(4-(2-amino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(316) 3-Z-[1-(4-(2-methylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (317) 3-Z-[1-(4-(2-ethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (318) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(319) 3-Z-[1-(4-(2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (320) 3-Z-[1-(4-(2-piperidin-1-yl-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(321) 3-Z-[1-(4-(2-acetylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(322) 3-Z-[1-(4-(3-amino-propyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(323) 3-Z-[1-(4-(3-dimethylamino-propyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(324) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(325) 3-Z-[1-(4-(N-ethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(326) 3-Z-[1-(4-(N-diethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(327) 3-Z-[1-(4-(N-dipropylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(328) 3-Z-[1-(4-(N-((N-ethyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(329) 3-Z-[1-(4-(N-((N-ethyl-N-propyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(330) 3-Z-[1-(4-(N-((N-methyl-N-propyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(331) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-ethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(332) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-propyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (333) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-butyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(334) 3-Z-[1-(4-(N-(2-amino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (335) 3-Z-[1-(4-(N-(2-diethylamino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(336) 3-Z-[1-(4-(N-acetyl-N-(2-aminoethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (337) 3-Z-[1-(4-(N-acetyl-N-(2-methylamino-ethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (338) 3-Z-[1-(4-(N-acetyl-N-(3-methylamino-propyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(339) 3-Z-[1-(4-(N-acetyl-N-(2-piperidin-1-yl-ethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (340) 3-Z-[1-(4-(N-acetyl-N-(aminocarbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(341) 3-Z-[1-(4-(N-acetyl-N-(piperidin-1-yl-carbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(342) 3-Z-[1-(4-(N-methyl-N-(aminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (343) 3-Z-[1-(4-(N-methyl-N-(methyaminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(344) 3-Z-[1-(4-(N-methyl-N-(dimethylaminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (345) 3-Z-[1-(4-(N-methyl-N-(piperidin-1-yl-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(346) 3-Z-[1-(4-(N-(2-ethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (347) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (348) 3-Z-[1-(4-(N-(2-pyrrolidin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(349) 3-Z-[1-(4-(N-(2-piperidin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (350) 3-Z-[1-(4-(N-(2-piperazin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(351) 3-Z-[1-(4-(N-(2-(4-morpholin-1-yl)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(352) 3-Z-[1-(4-(N-(ethylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (353) 3-Z-[1-(4-(N-(diethylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(354) 3-Z-[1-(4-(N-(pyrrolidin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (355) 3-Z-[1-(4-(N-(piperidin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(356) 3-Z-[1-(4-(N-(piperazin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (357) 3-Z-[1-(4-(N-((morpholin-4-yl)-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (358) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(359) 3-Z-[1-(4-(3-dimethylamino-propoxy)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (360) 3-Z-[1-(4-(aminocarbonylmethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(361) 3-Z-[1-(4-(2-aminocarbonyl-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(362) 3-Z-[1-(4-(pyridin-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (363) 3-Z-[1-(4-(pyridine-3-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(364) 3-Z-[1-(4((N-phenethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (365) 3-Z-[1-(4-(N-acetyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(366) 3-Z-[1-(4-(N-ethylcarbonyl-N-(dimethylaminocarbonyl-methyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (367) 3-Z-[1-(4-(N-methyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (368) 3-Z-[1-(4-carboxymethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(369) 3-Z-[1-(4-carbamoylmethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (370) 3-Z-[1-(4-dimethylcarbamoylmethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30 (371) 3-Z-[1-(4-tetrazol-5-yl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(372) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

5 (373) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(374) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

10 (375) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(376) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

15 (377) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

20 (378) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(379) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

25 (380) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(381) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

30

(382) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

5 (383) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(384) 3-Z-[1-(4-tetrazol-5-yl-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

10 (385) 3-Z-[1-(4-tetrazol-5-yl-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(386) 3-Z-[1-(4-tetrazol-5-yl-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(387) 3-Z-[1-(4-tetrazol-5-yl-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

15 (388) 3-Z-[1-(4-carboxy-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(389) 3-Z-[1-(4-carboxy-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(390) 3-Z-[1-(4-carboxy-anilino)-propylidene]-6-methoxy-carbonyl-2-indolinone

20 (391) 3-Z-[1-(4-carboxy-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(392) 3-Z-[1-(4-(N-benzyl-N-methyl-aminomethyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (393) 3-Z-[1-(4-(2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

30 (394) 3-Z-[1-(4-((benzo(1,3)dioxol-5-yl-methyl)-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(395) 3-Z-[1-(4-(N-phenethyl-N-methyl-aminomethyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (396) 3-Z-[1-(4-(N-(3,4-dimethoxy-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(397) 3-Z-[1-(4-(N-(4-Chloro-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (398) 3-Z-[1-(4-(N-(4-methylbenzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(399) 3-Z-[1-(4-(N-(4-fluoro-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (400) 3-Z-[1-(4-(N-(4-bromo-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (401) 3-Z-[1-(4-(N-(3-dimethylamino-propionyl)-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(402) 3-Z-[1-(4-(N-(4-dimethylamino-butyryl)-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (403) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-(2-dimethylamino-ethylsulphonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(404) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-(3-dimethylamino-propylsulphonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(405) 3-Z-[1-(4-((2-hydroxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (406) 3-Z-[1-(4-((2-methoxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(407) 3-Z-[1-(4-((2-dimethylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (408) 3-Z-[1-(4-((3-dimethylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(409) 3-Z-[1-(4-((N-tert.butoxycarbonyl-2-amino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (410) 3-Z-[1-(4-((N-tert.butoxycarbonyl-3-amino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (411) 3-Z-[1-(4-((2-amino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(412) 3-Z-[1-(4-((3-amino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (413) 3-Z-[1-(4-((2-acetylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(414) 3-Z-[1-(4-((3-acetylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(415) 3-Z-[1-(4-((2-methylsulphonylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (416) 3-Z-[1-(4-((3-methylsulphonylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(417) 3-Z-[1-(4-(N-(N-tert.butoxycarbonyl-2-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (418) 3-Z-[1-(4-(N-(2-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(419) 3-Z-[1-(4-(N-(2-acetyl-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (420) 3-Z-[1-(4-(N-(2-methylsulphonylamino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (421) 3-Z-[1-(4-(carboxymethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(422) 3-Z-[1-(4-(ethoxycarbonylmethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (423) 3-Z-[1-(4-(carbamoylemethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(424) 3-Z-[1-(4-(dimethylcarbamoylemethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(425) 3-Z-[1-(4-(methylcarbamoyl-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (426) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(427) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-nitro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (428) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-acetyl-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(429) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methylsulphonylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (430) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-cyano-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (431) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-hydroxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(432) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methoxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (433) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-ethoxycarbonyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(434) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-carboxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(435) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-carbamoyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(436) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-chloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(437) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-fluoro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(438) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-bromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(439) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(440) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-trifluoromethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(441) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3,5-dibromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(442) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3,5-dichloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(443) 3-Z-[1-(4-(dimethylaminomethyl)-3-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(444) 3-Z-[1-(4-(dimethylaminomethyl)-3-nitro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(445) 3-Z-[1-(4-(dimethylaminomethyl)-3-acetylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (446) 3-Z-[1-(4-(dimethylaminomethyl)-3-methylsulphonylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(447) 3-Z-[1-(4-(dimethylaminomethyl)-3-cyano-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (448) 3-Z-[1-(4-(dimethylaminomethyl)-3-hydroxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(449) 3-Z-[1-(4-(dimethylaminomethyl)-3-methoxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (450) 3-Z-[1-(4-(dimethylaminomethyl)-3-ethoxycarbonyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

20 (451) 3-Z-[1-(4-(dimethylaminomethyl)-3-carboxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(452) 3-Z-[1-(4-(dimethylaminomethyl)-3-carbamoyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (453) 3-Z-[1-(4-(dimethylaminomethyl)-3-chloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(454) 3-Z-[1-(4-(dimethylaminomethyl)-3-fluoro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

(455) 3-Z-[1-(4-(dimethylaminomethyl)-3-bromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

5 (456) 3-Z-[1-(4-(dimethylaminomethyl)-3-methyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(457) 3-Z-[1-(4-(dimethylaminomethyl)-3-trifluoromethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

10 (458) 3-Z-[1-(4-dimethylaminomethyl-3,5-dibromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(459) 3-Z-[1-(4-(dimethylaminomethyl)-3,5-dichloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

15 (460) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(2-hydroxy-ethoxy)-carbonyl]-2-indolinone

20 (461) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(ethoxycarbonyl-methoxy)-carbonyl]-2-indolinone

(462) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(carboxy-methoxy)-carbonyl]-2-indolinone

25 (463) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(carbamoylethoxy)-carbonyl]-2-indolinone

30 (464) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-hydroxy-ethoxy)-carbonyl]-2-indolinone

- (465) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(ethoxycarbonyl-methoxy)-carbonyl]-2-indolinone
- (466) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(carboxy-methoxy)-carbonyl]-2-indolinone
- (467) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(carbamoyl-methoxy)-carbonyl]-2-indolinone
- (468) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-methoxy-ethoxy)-carbonyl]-2-indolinone
- (469) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-dimethylamino-ethoxy)-carbonyl]-2-indolinone
- (470) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-(N-tert.butoxycarbonyl-amino)-ethoxy)-carbonyl]-2-indolinone
- (471) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-amino-ethoxy)-carbonyl]-2-indolinone
- (472) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2,2,2-trifluoroethoxy)-carbonyl]-2-indolinone
- (473) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (474) 3-Z-[1-(4-(N-(imidazo-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

- (475) 3-Z-[1-(4-(N-(phthalimido-2-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (476) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-
5 6-methoxycarbonyl-2-indolinone
- (477) 3-Z-[1-(4-(N-acetylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (478) 3-Z-[1-(4-(N-methylsulphonylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (479) 3-Z-[1-(4-(N-((N-(2-methoxyethyl)-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 15 (480) 3-Z-[1-(4-(N-((N-(2-dimethylaminoethyl)-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (481) 3-Z-[1-(4-(N-((di-(2-hydroxyethyl)-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 20 (482) 3-Z-[1-(4-(tert.butoxycarbonylmethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (483) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-methylene]-
25 6-methoxycarbonyl-2-indolinone
- (484) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-ethylidene]-
6-methoxycarbonyl-2-indolinone

30

(485) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(486) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-butylidene]-
5 6-methoxycarbonyl-2-indolinone

(487) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

10 (488) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(489) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

15 (490) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(491) 3-Z-[1-(4-tert.butylloxycarbonyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-
20 2-indolinone

(492) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

25 (493) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(494) 3-Z-[1-(4-(N-methyl-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

30

- (495) 3-Z-[1-(4-(imidazol-4-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- (496) 3-Z-[1-(4-((N-(dioxolan-2-yl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone
- 5 (497) 3-Z-[1-(4-(N-benzyl-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone
- 10 (498) 3-Z-[1-(4-(2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone
- (499) 3-Z-[1-(4-((benzo(1,3)dioxol-5-yl-methyl)-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone
- 15 (500) 3-Z-[1-(4-(N-phenethyl-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone
- (501) 3-Z-[1-(4-(N-(3,4-dimethoxy-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone
- 20 (502) 3-Z-[1-(4-(N-(4-Chloro-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone
- (503) 3-Z-[1-(4-(N-(4-methyl-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone
- 25 (504) 3-Z-[1-(4-(N-(4-fluoro-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

30

(505) 3-Z-[1-(4-(N-(4-bromo-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

5 (506) 3-Z-[1-(4-((N-(2-methoxy-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(507) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(2-amino-ethoxy)-carbonyl]-2-indolinone

10 (508) 3-Z-[1-(4-((N-(3-methylsulfonylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

Example 13

15 Dry ampoule containing 75 mg of active substance per 10 ml

Composition:

Active substance	75.0 mg
20 Mannitol	50.0 mg
water for injection	ad 10.0 ml

Preparation:

25 Active substance and mannitol are dissolved in water. After packaging the solution is freeze-dried. To produce the solution ready for use, the product is dissolved in water for injections.

Example 14

30 Dry ampoule containing 35 mg of active substance per 2 ml

Composition:

	Active substance	35.0 mg
	Mannitol	100.0 mg
5	water for injections	ad 2.0 ml

Preparation:

Active substance and mannitol are dissolved in water. After packaging, the solution is freeze-dried.

10

To produce the solution ready for use, the product is dissolved in water for injections.

Example 15

15 Tablet containing 50 mg of active substance

Composition:

	(1) Active substance	50.0 mg
20	(2) Lactose	98.0 mg
	(3) Maize starch	50.0 mg
	(4) Polyvinylpyrrolidone	15.0 mg
	(5) Magnesium stearate	<u>2.0 mg</u>
		215.0 mg

25

Preparation:

(1), (2) and (3) are mixed together and granulated with an aqueous solution of (4). (5) is added to the dried granulated material. From this mixture tablets are pressed, biplanar, faceted on both sides and with a dividing notch on one side.

30 Diameter of the tablets: 9 mm.

Example 16

Tablet containing 350 mg of active substance

5 Composition:

	(1) Active substance	350.0 mg
	(2) Lactose	136.0 mg
	(3) Maize starch	80.0 mg
10	(4) Polyvinylpyrrolidone	30.0 mg
	(5) Magnesium stearate	<u>4.0 mg</u>
		600.0 mg

Preparation:

- 15 (1), (2) and (3) are mixed together and granulated with an aqueous solution of (4). (5) is added to the dried granulated material. From this mixture tablets are pressed, biplanar, faceted on both sides and with a dividing notch on one side.

Diameter of the tablets: 12 mm.

20 Example 17

Capsules containing 50 mg of active substance

Composition:

25	(1) Active substance	50.0 mg
	(2) Dried maize starch	58.0 mg
	(3) Powdered lactose	50.0 mg
	(4) Magnesium stearate	<u>2.0 mg</u>
30		160.0 mg

Preparation:

(1) is triturated with (3). This trituration is added to the mixture of (2) and (4) with vigorous mixing.

- 5 This powder mixture is packed into size 3 hard gelatine capsules in a capsule filling machine.

Example 18

- 10 Capsules containing 350 mg of active substance
-

Composition:

- | | | |
|----|------------------------|---------------|
| | (1) Active substance | 350.0 mg |
| 15 | (2) Dried maize starch | 46.0 mg |
| | (3) Powdered lactose | 30.0 mg |
| | (4) Magnesium stearate | <u>4.0 mg</u> |
| | | 430.0 mg |

- 20 Preparation:

(1) is triturated with (3). This trituration is added to the mixture of (2) and (4) with vigorous mixing.

- 25 This powder mixture is packed into size 0 hard gelatine capsules in a capsule filling machine.

Example 19

Suppositories containing 100 mg of active substance

30

1 suppository contains:

Active substance	100.0 mg
Polyethyleneglycol (M.W. 1500)	600.0 mg
Polyethyleneglycol (M.W. 6000)	460.0 mg
Polyethylenesorbitan monostearate	<u>840.0 mg</u>
5	2,000.0 mg

Preparation:

The polyethyleneglycol is melted together with polyethylene sorbitan monostearate. At 40°C the ground active substance is homogeneously dispersed in the melt. It is cooled to 10 38°C and poured into slightly chilled suppository moulds.